

# Blow-up phenomena for scalar-flat metrics on manifolds with boundary

Sérgio de Moura Almaraz

## Abstract

Let  $(M^n, g)$  be a compact Riemannian manifold with boundary  $\partial M$ . This article is concerned with the set of scalar-flat metrics on  $M$  which are in the conformal class of  $g$  and have  $\partial M$  as a constant mean curvature hypersurface. We construct examples of metrics on the unit ball  $B^n$ , in dimensions  $n \geq 25$ , for which this set is noncompact. These manifolds have umbilic boundary, but they are not conformally equivalent to  $B^n$ .

## 1 Introduction

Let  $(M^n, g)$  be a compact Riemannian manifold with boundary  $\partial M$  and dimension  $n \geq 3$ . In 1992, J. Escobar addressed the question of finding a scalar-flat conformal metric  $\tilde{g} = u^{\frac{4}{n-2}}g$  which has  $\partial M$  as a constant mean curvature hypersurface. This problem was studied in [2], [9], [16], [17], [18], [27] and [28]. In analytical terms, it corresponds to the existence of a positive solution to the equations

$$\begin{cases} \Delta_g u - c_n R_g u = 0, & \text{in } M, \\ \frac{\partial u}{\partial \eta} - d_n \kappa_g u + K u^{\frac{n-2}{n-1}} = 0, & \text{on } \partial M, \end{cases} \quad (1.1)$$

for some constant  $K$ , where  $c_n = \frac{n-2}{4(n-1)}$  and  $d_n = \frac{n-2}{2}$ . Here,  $\Delta_g$  is the Laplace-Beltrami operator,  $R_g$  is the scalar curvature,  $\kappa_g$  is the mean curvature of  $\partial M$  and  $\eta$  is the inward unit normal vector to  $\partial M$ .

Escobar's question was motivated by the classical Yamabe problem, which consists of finding a conformal metric of constant scalar curvature on a given closed Riemannian manifold. This was completely solved after the works of H. Yamabe ([35]), N. Trudinger ([34]), T. Aubin ([4]) and R. Schoen ([30]). (See [22] and [32] for nice surveys on the issue.) Conformal metrics of constant scalar curvature and zero boundary mean curvature on the boundary were studied in [7], [15] (see also [3] and [20]).

The solutions to the equations (1.1) are the critical points of the functional

$$Q(u) = \frac{\int_M |du|_g^2 + c_n R_g u^2 dv_g + \int_{\partial M} d_n \kappa_g u^2 d\sigma_g}{\left( \int_{\partial M} |u|^{\frac{2(n-1)}{n-2}} d\sigma_g \right)^{\frac{n-2}{n-1}}},$$

where  $dv_g$  and  $d\sigma_g$  denote the volume forms of  $M$  and  $\partial M$ , respectively. In order to prove the existence of these solutions, Escobar introduced the conformally invariant Sobolev quotient

$$Q(M, \partial M) = \inf\{Q(u); u \in C^1(\bar{M}), u \neq 0 \text{ on } \partial M\}.$$

In this work we are interested in the question of whether the full set of solutions to (1.1) is compact. A necessary condition is that  $M$  is not conformally equivalent to the standard ball  $B^n$ . We point out that if the equations (1.1) have a solution  $u > 0$  with  $K$  positive (resp. zero and negative), then  $Q(M, \partial M)$  has to be positive (resp. zero and negative). If  $K < 0$ , the solution to the equations (1.1) is unique. If  $K = 0$ , the equations (1.1) become linear and the solutions are unique up to a multiplication by a positive constant. Hence, the only interesting case is the one when  $K > 0$ .

The problem of compactness of solutions to the equations (1.1) was studied by V. Felli and M. Ould Ahmedou in the conformally flat case with umbilic boundary ([18]) and in the three-dimensional case with umbilic boundary ([19]). In [1], the author proved compactness for dimensions  $n \geq 7$  under a generic condition. Other compactness results for similar equations were obtained by Z. Djadli, A. Malchiodi and M. Ould Ahmedou in [11, 12], by Z. Han and Y. Li in [20] and by M. Ould Ahmedou in [29].

In the case of manifolds without boundary, the question of compactness of the full set of solutions to the Yamabe equation was first raised by R. Schoen in a topics course at Stanford University in 1988. A necessary condition is that the manifold  $M^n$  is not conformally equivalent to the sphere  $S^n$ . This problem was studied in [13], [14], [23], [24], [25], [26], [31] and [33] and was completely solved in a series of three papers: [6], [8] and [21]. In [6], S. Brendle discovered the first smooth counterexamples for dimensions  $n \geq 52$  (see [5] for nonsmooth examples). In [21], M. Khuri, F. Marques and R. Schoen proved compactness for dimensions  $3 \leq n \leq 24$ . Finally, in [8], Brendle and Marques extended the counterexamples of [6] to the remaining dimensions  $25 \leq n \leq 51$ .

It is expected that, as in the case of manifolds without boundary, there should be a critical dimension  $n_0$  such that compactness in the case of manifolds with boundary holds for  $n < n_0$  and fails for  $n \geq n_0$ . In this work we partially answer this question by showing that compactness fails for dimensions  $n \geq 25$ . More precisely we prove:

**Main Theorem.** *Let  $n \geq 25$ . Then there exists a smooth Riemannian metric  $g$  on  $B^n$  and a sequence of positive smooth functions  $\{v_\nu\}_{\nu=1}^\infty$  with the following properties:*

- (i)  $g$  is not conformally flat;
- (ii)  $\partial B^n$  is umbilic with respect to the induced metric by  $g$ ;
- (iii) for all  $\nu$ ,  $v_\nu$  is a solution to the equations (1.1) with a constant  $K > 0$  and  $M = B^n$ ;
- (iv)  $Q(v_\nu) < Q(B^n, \partial B)$  for all  $\nu$ ;
- (v)  $\sup_{\partial B^n} v_\nu \rightarrow \infty$  as  $\nu \rightarrow \infty$ .

In order to prove the Main Theorem, we follow the program adopted in [6] and [8]. In Section 2, we show that the problem can be reduced to finding critical points of a certain function  $\mathcal{F}_g(\xi, \epsilon)$ , where  $\xi$  is a vector in  $\mathbb{R}^{n-1}$  and  $\epsilon$  is a positive real number. In Section 3, we show that the function  $\mathcal{F}_g(\xi, \epsilon)$  can be approximated by an auxiliary function  $F(\xi, \epsilon)$ . In Section 4, we prove that the function  $F(\xi, \epsilon)$  has a strict local minimum point. The cases  $n \geq 53$  and  $25 \leq n \leq 52$  are handled separately in Subsections 4.1 and 4.2 respectively. Finally, in Section 5, we use a perturbation argument to construct critical points of the function  $\mathcal{F}_g(\xi, \epsilon)$  and prove the non-compactness theorem.

**Notation.** Throughout this work we will make use of the index notation for tensors. We will adopt the summation convention whenever confusion is not possible and use indices  $1 \leq i, j, k, l, m, p, q, r, s \leq n-1$  and  $1 \leq a, b, c, d \leq n$ . We also define constants  $c_n = \frac{n-2}{4(n-1)}$  and  $d_n = \frac{n-2}{2}$ .

We will denote by  $\Delta_g$  the Laplace-Beltrami operator. The volume forms of  $M$  and  $\partial M$  will be denoted by  $dv_g$  and  $d\sigma_g$ , respectively. By  $\eta$  we will denote the inward unit normal vector to  $\partial M$ . The scalar curvature will be denoted by  $R_g$ , the second fundamental form of  $\partial M$  by  $\pi_{kl}$  and the mean curvature,  $\frac{1}{n-1} \text{tr}(\pi_{kl})$ , by  $\kappa_g$ .

By  $\mathbb{R}_+^n$  we will denote the half-space  $\{x = (x_1, \dots, x_n) \in \mathbb{R}^n; x_n \geq 0\}$ . If  $x \in \mathbb{R}_+^n$  we set  $\bar{x} = (x_1, \dots, x_{n-1}, 0) \in \partial\mathbb{R}_+^n \cong \mathbb{R}^{n-1}$ . For any  $x_0 \in \mathbb{R}_+^n$  we set  $B_r^+(x_0) = \{x \in \mathbb{R}_+^n; |x - x_0| < r\}$ . The  $n$ -dimensional sphere of radius  $r$  in  $\mathbb{R}^{n+1}$  will be denoted by  $S_r^n$  and  $\sigma_n$  will denote the area of the  $n$ -dimensional unit sphere  $S_1^n$ .

**Acknowledgements.** I would like to thank Prof. Fernando C. Marques for his comments and interest in this work.

## 2 Lyapunov-Schmidt reduction

Given a pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$  we set

$$u_{(\xi, \epsilon)}(x) = \left( \frac{\epsilon}{(\epsilon + x_n)^2 + |\bar{x} - \xi|^2} \right)^{\frac{n-2}{2}}, \quad \text{for } x \in \mathbb{R}_+^n.$$

Observe that  $u_{(\xi, \epsilon)}$  satisfies

$$\begin{cases} \Delta u_{(\xi, \epsilon)} = 0, & \text{in } \mathbb{R}_+^n, \\ \frac{\partial}{\partial x_n} u_{(\xi, \epsilon)} + (n-2)u_{(\xi, \epsilon)}^{\frac{n-2}{n}} = 0, & \text{on } \partial\mathbb{R}_+^n, \end{cases} \quad (2.1)$$

and

$$\int_{\partial\mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{2(n-1)}{n-2}} = \left( \frac{Q(B^n, \partial B)}{n-2} \right)^{n-1}. \quad (2.2)$$

Let us define

$$\phi_{(\xi, \epsilon, n)}(x) = \left( \frac{\epsilon}{(\epsilon + x_n)^2 + |\bar{x} - \xi|^2} \right)^{\frac{n}{2}} \frac{\epsilon^2 - x_n^2 - |\bar{x} - \xi|^2}{(\epsilon + x_n)^2 + |\bar{x} - \xi|^2}$$

and

$$\phi_{(\xi, \epsilon, k)}(x) = \left( \frac{\epsilon}{(\epsilon + x_n)^2 + |\bar{x} - \xi|^2} \right)^{\frac{n}{2}} \frac{2\epsilon(x_k - \xi_k)}{(\epsilon + x_n)^2 + |\bar{x} - \xi|^2}$$

for  $x \in \mathbb{R}_+^n$  and  $k = 1, \dots, n-1$ . Observe that

$$\phi_{(\xi, \epsilon, n)}(x) \cdot ((\epsilon + x_n)^2 + |\bar{x} - \xi|^2) = -\frac{2\epsilon^2}{n-2} \frac{\partial}{\partial \epsilon} u_{(\xi, \epsilon)}(x),$$

$$\phi_{(\xi, \epsilon, k)}(x) \cdot ((\epsilon + x_n)^2 + |\bar{x} - \xi|^2) = \frac{2\epsilon^2}{n-2} \frac{\partial}{\partial \xi_k} u_{(\xi, \epsilon)}(x),$$

for  $k = 1, \dots, n-1$ , and that  $\|\phi_{(\xi, \epsilon, a)}\|_{L^{\frac{2(n-1)}{n}}(\partial\mathbb{R}_+^n)}$  is independent of  $(\epsilon, \xi) \in \mathbb{R}^{n-1} \times (0, \infty)$ , for any  $a = 1, \dots, n$ .

We also set

$$\Sigma = \left\{ w \in L^{\frac{2n}{n-2}}(\mathbb{R}_+^n) \cap L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n) \cap H_{loc}^1(\mathbb{R}_+^n); \int_{\mathbb{R}_+^n} |dw|^2 < \infty \right\},$$

$$\Sigma_{(\xi, \epsilon)} = \left\{ w \in \Sigma; \int_{\partial\mathbb{R}_+^n} \phi_{(\xi, \epsilon, a)} w = 0, a = 1, \dots, n \right\}$$

and  $\|w\|_{\Sigma} = \left( \int_{\mathbb{R}_+^n} |dw|^2 \right)^{\frac{1}{2}}$  for  $w \in \Sigma$ . Observe that  $u_{(\xi, \epsilon)} \in \Sigma_{(\xi, \epsilon)}$  for each  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$ . By Sobolev's inequality, there exists  $K = K(n) > 0$  such that

$$\left( \int_{\mathbb{R}_+^n} |w|^{\frac{2n}{n-2}} \right)^{\frac{n-2}{n}} + \left( \int_{\partial\mathbb{R}_+^n} |w|^{\frac{2(n-1)}{n-2}} \right)^{\frac{n-2}{n-1}} \leq K \int_{\mathbb{R}_+^n} |dw|^2 \quad (2.3)$$

for all  $w \in \Sigma$ .

In what follows in this section we are going to find, for each pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$ , a function  $v_{(\xi, \epsilon)} \in \Sigma$  which is an approximate weak solution to a Yamabe-type problem (1.1) on  $\mathbb{R}_+^n$ . Then we will show that  $v_{(\xi, \epsilon)}$  is in fact a classical solution to this problem whenever  $(\xi, \epsilon)$  is a critical point of a certain energy function defined on  $\mathbb{R}^{n-1} \times (0, \infty)$ .

The following result is Proposition 26 of [6] and will be used throughout this work:

**Lemma 2.1.** *Suppose that we express the Riemannian metric  $g$  as  $g = \exp(h)$ , where  $h$  is a trace-free symmetric two-tensor defined on  $\mathbb{R}_+^n$  and satisfying  $|h(x)| \leq 1$  for any  $x \in \mathbb{R}_+^n$ . Then there exists  $C = C(n) > 0$  such that*

$$\left| R_g - \left\{ \partial_a \partial_b h_{ab} - \partial_a (h_{ac} \partial_b h_{bc}) + \frac{1}{2} \partial_a h_{ac} \partial_b h_{bc} - \frac{1}{4} \partial_c h_{ab} \partial_c h_{ab} \right\} \right| \leq C|h|^2 |\partial^2 h| + C|h| |\partial h|^2.$$

**Notation.** In this section we suppose that  $g$  is a Riemannian metric on  $\mathbb{R}_+^n$  expressed as  $g = \exp(h)$ , where  $h$  is a trace-free symmetric two-tensor satisfying  $h(x) = 0$  for any  $|x| \geq 1$ .

**Proposition 2.2.** *If  $|h(x)| + |\partial h(x)| + |\partial^2 h(x)| \leq \alpha \leq 1$  for any  $x \in \mathbb{R}_+^n$ , then there exists  $C = C(n) > 0$  such that*

$$\left\| \Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)} \right\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} + \left\| d_n \kappa_g u_{(\xi, \epsilon)} \right\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)} \leq C\alpha$$

for all pairs  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$ .

*Proof.* It follows from the pointwise estimates

$$\left| \Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)} \right| \leq C \left\{ |h| |\partial^2 u_{(\xi, \epsilon)}| + |\partial h| |\partial u_{(\xi, \epsilon)}| + (|\partial^2 h| + |\partial h|^2) |u_{(\xi, \epsilon)}| \right\}$$

and  $|d_n \kappa_g u_{(\xi, \epsilon)}| \leq C |\partial h| |u_{(\xi, \epsilon)}|$  that

$$\begin{aligned} & \left\| \Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)} \right\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \\ & \leq C \left\{ \|h\|_{L^\infty(\mathbb{R}_+^n)} \|\partial^2 u_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} + \|\partial h\|_{L^n(\mathbb{R}_+^n)} \|\partial u_{(\xi, \epsilon)}\|_{L^2(\mathbb{R}_+^n)} \right\} \\ & \quad + C \left\{ \|\partial^2 h\|_{L^{\frac{n}{2}}(\mathbb{R}_+^n)} + \|\partial h\|_{L^n(\mathbb{R}_+^n)}^2 \right\} \|u\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)} \end{aligned}$$

and

$$\|d_n \kappa_g u_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)} \leq C \|\partial h\|_{L^{n-1}(\partial \mathbb{R}_+^n)} \|u_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)}.$$

From this the result follows.  $\square$

**Lemma 2.3.** *Let  $B^n = B_{1/2}^n(0, \dots, 0, -\frac{1}{2}) \subset \mathbb{R}^n$  be the ball with radius  $\frac{1}{2}$  and center  $(0, \dots, 0, -\frac{1}{2})$ . Let  $z_1, \dots, z_n$  be the coordinates of  $B^n$  taken with center  $(0, \dots, 0, -\frac{1}{2})$ . For each pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$ , the expression*

$$C_{(\xi, \epsilon)}(x) = \frac{\epsilon(x_1 - \xi_1, \dots, x_{n-1} - \xi_{n-1}, x_n + \epsilon)}{|\bar{x} - \xi|^2 + (x_n + \epsilon)^2} + (0, \dots, 0, -1)$$

defines a conformal equivalence

$$C_{(\xi, \epsilon)} : \mathbb{R}_+^n \rightarrow B^n \setminus \{(0, \dots, 0, -1)\}$$

that satisfies  $C_{(\xi, \epsilon)}^* \delta_{B^n} = u_{(\xi, \epsilon)}^{\frac{4}{n-2}} \delta$ , where  $\delta_{B^n}$  is the Euclidean metric on  $B^n$  and  $\delta$  is the Euclidean metric on  $\mathbb{R}_+^n$ . For any smooth function  $f$  on  $\mathbb{R}_+^n$ , we have

$$\Delta_{B^n} \tilde{u}_{(\xi, \epsilon)} = u_{(\xi, \epsilon)}^{-\frac{n+2}{n-2}} \Delta f \tag{2.4}$$

and

$$\frac{\partial}{\partial \eta} \tilde{u}_{(\xi, \epsilon)} - (n-2) \tilde{u}_{(\xi, \epsilon)} = u_{(\xi, \epsilon)}^{-\frac{n}{n-2}} \frac{\partial f}{\partial x_n}, \tag{2.5}$$

where  $\tilde{u}_{(\xi,\epsilon)} = (fu_{(\xi,\epsilon)}^{-1}) \circ C_{(\xi,\epsilon)}^{-1}$ . Moreover,

$$z_n \circ C_{(\xi,\epsilon)} = -\frac{\epsilon}{n-2} u_{(\xi,\epsilon)}^{-1} \frac{\partial}{\partial \epsilon} u_{(\xi,\epsilon)} = \frac{1}{2} u_{(\xi,\epsilon)}^{-\frac{n}{n-2}} \phi_{(\xi,\epsilon,n)} \quad (2.6)$$

and

$$z_k \circ C_{(\xi,\epsilon)} = \frac{\epsilon}{n-2} u_{(\xi,\epsilon)}^{-1} \frac{\partial}{\partial \xi_k} u_{(\xi,\epsilon)} = \frac{1}{2} u_{(\xi,\epsilon)}^{-\frac{n}{n-2}} \phi_{(\xi,\epsilon,k)}, \quad k = 1, \dots, n-1. \quad (2.7)$$

*Proof.* These are direct computations. The assertions (2.4) and (2.5) follow from the following properties of the conformal operators  $L_g = \Delta_g - c_n R_g$  and  $B_g = \frac{\partial}{\partial \eta} - d_n \kappa_g$ :

$$L_{u^{\frac{4}{n-2}}g}(fu^{-1}) = u^{-\frac{n+2}{n-2}} L_g f \quad \text{and} \quad B_{u^{\frac{4}{n-2}}g}(fu^{-1}) = u^{-\frac{n}{n-2}} B_g f. \quad (2.8)$$

□

**Lemma 2.4.** *There exists  $\theta = \theta(n) > 0$  such that*

$$\int_{B^n} |dw|^2 - 2 \int_{\partial B^n} w^2 - 2\theta \left( \int_{B^n} |dw|^2 + (n-2) \int_{\partial B^n} w^2 \right) + \frac{4}{\theta} \left( \int_{\partial B^n} w \right)^2 \geq 0$$

for any  $w \in H^1(B^n)$  such that  $w \perp_{L^2(\partial B^n)} \{z_1, \dots, z_n\}$ . Here, we are following the notations of Lemma 2.3.

*Proof.* First we fix  $0 \neq w \in H^1(B^n)$  such that  $w \perp_{L^2(\partial B^n)} \{1, z_1, \dots, z_n\}$ . Since

$$\inf \left\{ \frac{\int_{B^n} |d\psi|^2}{\int_{\partial B^n} \psi^2}, \text{ such that } \psi \in H^1(B^n), \psi \neq 0 \text{ on } \partial B^n \text{ and } \psi \perp_{L^2(\partial B^n)} 1 \right\} = 2$$

and this infimum is realized only by the functions  $z_1, \dots, z_n$ , we see that

$$\int_{B^n} |dw|^2 - 2 \int_{\partial B^n} w^2 > 0.$$

Hence,

$$\int_{B^n} |dw|^2 - 2 \int_{\partial B^n} w^2 \geq 2\theta \left( \int_{B^n} |dw|^2 + (n-2) \int_{\partial B^n} w^2 \right) \quad (2.9)$$

holds for any  $\theta > 0$  satisfying

$$\theta \leq \theta(w) = \frac{1}{2} \frac{\int_{B^n} |dw|^2 - 2 \int_{\partial B^n} w^2}{\int_{B^n} |dw|^2 + (n-2) \int_{\partial B^n} w^2}$$

and the equality is realized by  $\theta = \theta(w)$ .

We claim that there exists  $\theta_0 > 0$  such that  $\theta(w) \geq \theta_0$  for any  $w \in H^1(B^n)$  satisfying  $w \perp_{L^2(\partial B^n)} \{1, z_1, \dots, z_n\}$ . Suppose by contradiction this is not true. Thus

we can choose a sequence  $\{w_j\}_{j=1}^\infty \subset H^1(B^n)$  such that  $w_j \perp_{L^2(\partial B^n)} \{1, z_1, \dots, z_n\}$  and  $\theta(w_j) \rightarrow 0$  as  $j \rightarrow \infty$ . Hence

$$\int_{B^n} |dw_j|^2 - 2 \int_{\partial B^n} w_j^2 = 2\theta(w_j) \left( \int_{B^n} |dw_j|^2 + (n-2) \int_{\partial B^n} w_j^2 \right)$$

holds and we can assume that  $\int_{B^n} |dw_j|^2 = 1$  for any  $j$ . Thus,  $\int_{\partial B^n} w_j^2 \leq \frac{1}{2}$  for all  $j$  and we can suppose that  $w_j \rightarrow w_0$  in  $H^1(B^n)$  for some  $w_0$ . Since  $H^1(B^n)$  is compactly imbedded in  $L^2(\partial B^n)$ , we know that  $w_0 \perp_{L^2(\partial B^n)} \{1, z_1, \dots, z_n\}$ . Let us first assume that  $w_0 \not\equiv 0$ . We set

$$\beta = \int_{B^n} |dw_0|^2 - 2 \int_{\partial B^n} w_0^2 > 0.$$

Since  $\liminf_{i \rightarrow \infty} \int_{B^n} |dw_j|^2 \geq \int_{B^n} |dw_0|^2$  and  $\lim_{i \rightarrow \infty} \int_{\partial B^n} w_j^2 = \int_{\partial B^n} w_0^2$ , we can assume that

$$\int_{B^n} |dw_j|^2 - 2 \int_{\partial B^n} w_j^2 \geq \frac{\beta}{2} \quad \text{for all } j.$$

On the other hand,

$$\frac{\beta}{n} \left\{ \int_{B^n} |dw_j|^2 + (n-2) \int_{\partial B^n} w_j^2 \right\} \leq \frac{\beta}{2},$$

since  $\int_{B^n} |dw_j|^2 = 1$  and  $\int_{\partial B^n} w_j^2 \leq \frac{1}{2}$ . Hence,

$$\begin{aligned} 2\theta(w_j) \left( \int_{B^n} |dw_j|^2 + (n-2) \int_{\partial B^n} w_j^2 \right) &= \int_{B^n} |dw_j|^2 - 2 \int_{\partial B^n} w_j^2 \\ &\geq \frac{\beta}{n} \left( \int_{B^n} |dw_j|^2 + (n-2) \int_{\partial B^n} w_j^2 \right). \end{aligned}$$

which implies that  $2\theta(w_j) \geq \frac{\beta}{n}$  for all  $j$  and contradicts the fact that  $\theta(w_j) \rightarrow 0$ .

Thus we must have  $w_0 \equiv 0$ , which implies that  $\int_{\partial B^n} w_j^2 \rightarrow 0$  as  $j \rightarrow \infty$ . Then, if we set  $\tilde{w}_j = \left( \int_{\partial B^n} w_j^2 \right)^{-\frac{1}{2}} w_j$ , we have  $\tilde{w}_j \rightarrow \tilde{w}_0$  in  $H^1(B^n)$ , for some  $\tilde{w}_0$ . Moreover,

$$0 = \lim_{j \rightarrow \infty} \int_{B^n} |d\tilde{w}_j|^2 \geq \int_{B^n} |d\tilde{w}_0|^2$$

and

$$\int_{\partial B^n} \tilde{w}_j^2 = 1 = \int_{\partial B^n} \tilde{w}_0^2.$$

From this we conclude that  $\tilde{w}_0 \equiv \text{const} \neq 0$ , which contradicts the fact that  $\tilde{w}_0 \perp_{L^2(\partial B^n)} 1$ . This proves that there exists  $\theta_0 > 0$  such that  $\theta(w) \geq \theta_0$  for any  $w \in H^1(B^n)$  satisfying  $w \perp_{L^2(\partial B^n)} \{1, z_1, \dots, z_n\}$ . In particular, (2.9) holds, with  $\theta = \theta_0$ , for any  $w \in H^1(B^n)$  satisfying  $w \perp_{L^2(\partial B^n)} \{1, z_1, \dots, z_n\}$ .

Now, let  $w \in H^1(B^n)$  satisfy  $w \perp_{L^2(\partial B^n)} \{z_1, \dots, z_n\}$ . We write  $w = w_1 + b$  where  $b$  is a constant and  $w_1 \perp_{L^2(\partial B^n)} 1$ . Then we have

$$\begin{aligned} & \int_{B^n} |dw|^2 - 2 \int_{\partial B^n} w^2 - 2\theta_0 \left( \int_{B^n} |dw|^2 + (n-2) \int_{\partial B^n} w^2 \right) + \frac{4}{\theta_0} \left( \int_{\partial B^n} w \right)^2 \\ &= \int_{B^n} |dw_1|^2 - 2 \int_{\partial B^n} w_1^2 - 2\theta_0 \left( \int_{B^n} |dw_1|^2 + (n-2) \int_{\partial B^n} w_1^2 \right) \\ &\quad - 2(1 + (n-2)\theta_0) \int_{\partial B^n} b^2 + \frac{4}{\theta_0} \left( \int_{\partial B^n} b \right)^2 \\ &\geq \left( \frac{4}{\theta_0} - 2 - 2(n-2)\theta_0 \right) \int_{\partial B^n} b^2 \end{aligned}$$

Choosing  $\theta_0$  smaller if necessary, we can suppose that  $\frac{4}{\theta_0} - 2 - 2(n-2)\theta_0 > 0$  and the result follows.  $\square$

**Proposition 2.5.** *There exists  $\theta = \theta(n) > 0$  such that*

$$\int_{\mathbb{R}_+^n} |dw|^2 - n \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2 \geq 2\theta |w|_\Sigma^2 - \frac{4}{\theta} \left( \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{n}{n-2}} w \right)^2$$

for all  $w \in \Sigma_{(\xi, \epsilon)}$  and any pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$ .

*Proof.* Let  $w \in \Sigma_{(\xi, \epsilon)}$  and set  $\bar{w} = (wu_{(\xi, \epsilon)}^{-1}) \circ C_{(\xi, \epsilon)}^{-1}$ . Using the fact that  $C_0^\infty(\mathbb{R}_+^n)$  is dense in  $\Sigma$  with respect to the norms  $\|\cdot\|_\Sigma$ ,  $\|\cdot\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)}$  and  $\|\cdot\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)}$ , it is easy to see that we can assume that  $\bar{w} \in H^1(B^n)$ . It follows from the expressions (2.6) and (2.7) that

$$\int_{\partial B^n} \bar{w} z_n = \frac{1}{2} \int_{\partial \mathbb{R}_+^n} w \phi_{(\xi, \epsilon, n)} = 0$$

and

$$\int_{\partial B^n} \bar{w} z_k = \frac{1}{2} \int_{\partial \mathbb{R}_+^n} w \phi_{(\xi, \epsilon, k)} = 0, \quad k = 1, \dots, n-1.$$

Then, according to Lemma 2.4, we have

$$\int_{B^n} |d\bar{w}|^2 - 2 \int_{\partial B^n} \bar{w}^2 - 2\theta \left( \int_{B^n} |d\bar{w}|^2 + (n-2) \int_{\partial B^n} \bar{w}^2 \right) + \frac{4}{\theta} \left( \int_{\partial B^n} \bar{w} \right)^2 \geq 0. \quad (2.10)$$

Hence, using the formulas (2.4) and (2.5), we easily see that

$$\begin{aligned} & \int_{B^n} |d\bar{w}|^2 - 2 \int_{\partial B^n} \bar{w}^2 = \int_{\mathbb{R}_+^n} |dw|^2 - n \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2, \\ & \int_{B^n} |d\bar{w}|^2 + (n-2) \int_{\partial B^n} \bar{w}^2 = \int_{\mathbb{R}_+^n} |dw|^2 \quad \text{and} \quad \int_{\partial B^n} \bar{w} = \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{n}{n-2}} w. \end{aligned}$$

Now the result follows from substituting these last three equations in (2.10).  $\square$



**Corollary 2.6.** *Let  $K$  be as in (2.3) and  $\theta$  be as in Proposition 2.5. Then there exists  $0 < \alpha_0 = \alpha_0(n) \leq 1$  such that, whenever  $|h(x)| + |\partial h(x)| + |\partial^2 h(x)| \leq \alpha_0$  for all  $x \in \mathbb{R}_+^n$ , we have*

$$\left( \int_{\mathbb{R}_+^n} |w|^{\frac{2n}{n-2}} \right)^{\frac{n-2}{n}} + \left( \int_{\partial \mathbb{R}_+^n} |w|^{\frac{2(n-1)}{n-2}} \right)^{\frac{n-2}{n-1}} \leq 2K \int_{\mathbb{R}_+^n} (|dw|_g^2 + c_n R_g w^2) + 2K \int_{\partial \mathbb{R}_+^n} d_n \kappa_g w^2 \quad (2.11)$$

for all  $w \in \Sigma$  and

$$\int_{\mathbb{R}_+^n} (|dw|_g^2 + c_n R_g w^2) + \int_{\partial \mathbb{R}_+^n} (d_n \kappa_g w^2 - nu_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2) \geq \frac{\theta}{2} \|w\|_\Sigma^2 - \frac{1}{\theta} A(w)^2 \quad (2.12)$$

for all  $w \in \Sigma_{(\xi, \epsilon)}$  and any pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$ . Here,

$$A(w) = \int_{\mathbb{R}_+^n} (\Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)}) w + \int_{\partial \mathbb{R}_+^n} (-d_n \kappa_g u_{(\xi, \epsilon)} + 2u_{(\xi, \epsilon)}^{\frac{n}{n-2}}) w.$$

*Proof.* Let us first prove the estimate (2.12). Observe that

$$\int_{\mathbb{R}_+^n} (\Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)}) w \geq - \|\Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \|w\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)}$$

and

$$\begin{aligned} & \int_{\partial \mathbb{R}_+^n} (-d_n \kappa_g u_{(\xi, \epsilon)} + 2u_{(\xi, \epsilon)}^{\frac{n}{n-2}}) w \\ & \geq - \|d_n \kappa_g u_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)} \|w\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)} + 2 \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{n}{n-2}} w \end{aligned}$$

Hence, by Proposition 2.2 and inequality (2.3) we have

$$A(w) \geq -C\alpha_0 \|w\|_\Sigma + 2 \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{n}{n-2}} w.$$

Choosing  $\alpha_0$  small this implies

$$A(w)^2 \geq 4 \left( \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{n}{n-2}} w \right)^2 - \theta^2 \|w\|_\Sigma^2$$

which, together with Proposition 2.5, gives

$$\int_{\mathbb{R}_+^n} |dw|^2 - \int_{\partial \mathbb{R}_+^n} nu_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2 \geq \theta \|w\|_\Sigma^2 - \frac{1}{\theta} A(w)^2. \quad (2.13)$$

On the other hand,

$$\begin{aligned} & \int_{\mathbb{R}_+^n} (|dw|_g^2 + c_n R_g w^2) + \int_{\partial \mathbb{R}_+^n} (d_n \kappa_g w^2 - nu_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2) \\ & = \int_{\mathbb{R}_+^n} |dw|^2 - \int_{\partial \mathbb{R}_+^n} nu_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2 + \int_{\mathbb{R}_+^n} \{(g^{ij} - \delta^{ij}) \partial_i w \partial_j w + c_n R_g w^2\} + \int_{\partial \mathbb{R}_+^n} d_n \kappa_g w^2. \end{aligned}$$

The fact that  $h(x) = 0$  for  $|x| \geq 1$  and (2.3) imply that

$$\begin{aligned} \int_{\mathbb{R}_+^n} \left\{ (g^{ij} - \delta^{ij}) \partial_i w \partial_j w + c_n R_g w^2 \right\} + \int_{\partial \mathbb{R}_+^n} d_n \kappa_g w^2 & \quad (2.14) \\ & \leq C \alpha_0 \|w\|_{\Sigma}^2 + C \alpha_0 \|w\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)}^2 + C \alpha_0 \|w\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)}^2 \\ & \leq C \alpha_0 (1 + K) \|w\|_{\Sigma}^2. \end{aligned}$$

Hence,

$$\begin{aligned} \int_{\mathbb{R}_+^n} (|dw|_g^2 + c_n R_g w^2) + \int_{\partial \mathbb{R}_+^n} (d_n \kappa_g w^2 - n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2) & \quad (2.15) \\ & \geq \int_{\mathbb{R}_+^n} |dw|^2 - \int_{\partial \mathbb{R}_+^n} n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2 - C \alpha_0 (1 + K) \|w\|_{\Sigma}^2. \end{aligned}$$

Now the result follows from the inequalities (2.13) and (2.15), choosing  $\alpha_0$  small. The estimate (2.11) follows easily from the inequalities (2.3) and (2.14).  $\square$

**Proposition 2.7.** *Suppose that  $|h(x)| + |\partial h(x)| + |\partial^2 h(x)| \leq \alpha_0$  for all  $x \in \mathbb{R}_+^n$ , where  $\alpha_0$  is the constant obtained in Corollary 2.6. Given any pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$  and functions  $f \in L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)$  and  $\bar{f} \in L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)$  there exists a unique  $w \in \Sigma_{(\xi, \epsilon)}$  such that*

$$\int_{\mathbb{R}_+^n} (\langle dw, d\psi \rangle_g + c_n R_g w \psi) + \int_{\partial \mathbb{R}_+^n} (d_n \kappa_g w \psi - n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w \psi) = \int_{\mathbb{R}_+^n} f \psi + \int_{\partial \mathbb{R}_+^n} \bar{f} \psi \quad (2.16)$$

for all  $\psi \in \Sigma_{(\xi, \epsilon)}$ . Moreover, there exists  $C = C(n) > 0$  such that

$$\|w\|_{\Sigma} \leq C \|f\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} + C \|\bar{f}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)}.$$

*Proof.* Let us first prove the existence part. Following the notations of Corollary 2.6, we define the functional

$$T(w) = \int_{\mathbb{R}_+^n} (|dw|_g^2 + c_n R_g w^2 - 2fw) + \int_{\partial \mathbb{R}_+^n} (d_n \kappa_g w^2 - n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2 - 2\bar{f}w) + \frac{1}{\theta} A(w)^2$$

for  $w \in \Sigma_{(\xi, \epsilon)}$ . Hence

$$\begin{aligned} dT_w(\psi) &= 2 \int_{\mathbb{R}_+^n} (\langle dw, d\psi \rangle_g + c_n R_g w \psi - f\psi) + 2 \int_{\partial \mathbb{R}_+^n} (d_n \kappa_g w \psi - n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w \psi - \bar{f}\psi) \\ &\quad + \frac{2}{\theta} A(w) A(\psi). \end{aligned}$$

It follows from the identity (2.12) that

$$\begin{aligned} T(w) &\geq \frac{\theta}{2} \|w\|_{\Sigma}^2 - 2 \int_{\mathbb{R}_+^n} f w - 2 \int_{\partial \mathbb{R}_+^n} \bar{f} w \\ &\geq \frac{\theta}{2} \|w\|_{\Sigma}^2 - 2 \|f\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \|w\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)} - 2 \|\bar{f}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)} \|w\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)} \\ &\geq \frac{\theta}{4} \|w\|_{\Sigma}^2 - C \left( \|f\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)}^2 + \|\bar{f}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)}^2 \right) \end{aligned}$$

where in the last inequality we used the estimate (2.3). So,  $T$  is bounded below and by a standard argument we can find a minimizer  $w_0$  for  $T$  over all functions in  $\Sigma_{(\xi, \epsilon)}$ . Now, integrating by parts we see that

$$\int_{\mathbb{R}_+^n} \left( \langle du_{(\xi, \epsilon)}, d\psi \rangle_g + c_n R_g u_{(\xi, \epsilon)} \psi \right) + \int_{\partial \mathbb{R}_+^n} \left( d_n \kappa_g u_{(\xi, \epsilon)} \psi - nu_{(\xi, \epsilon)}^{\frac{n}{n-2}} \psi \right) = -A(\psi),$$

holds for all  $\psi \in C_0^\infty(\mathbb{R}_+^n)$ . Since this space is dense in  $\Sigma$  with respect to the norms  $\|\cdot\|_\Sigma$ ,  $\|\cdot\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)}$  and  $\|\cdot\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)}$ , this identity holds for all  $\psi \in \Sigma$ . Hence, the function  $w = w_0 - \frac{1}{\theta} A(w_0) u_{(\xi, \epsilon)}$  satisfies (2.16) for all  $\psi \in \Sigma_{(\xi, \epsilon)}$ , proving the existence part.

In order to prove the uniqueness part, suppose that  $w \in \Sigma_{(\xi, \epsilon)}$  satisfies (2.16) for all  $\psi \in \Sigma_{(\xi, \epsilon)}$ . In particular,

$$\int_{\mathbb{R}_+^n} \left( |dw|_g^2 + c_n R_g w^2 \right) + \int_{\partial \mathbb{R}_+^n} \left( d_n \kappa_g w^2 - nu_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2 \right) = \int_{\mathbb{R}_+^n} f w + \int_{\partial \mathbb{R}_+^n} \bar{f} w$$

and

$$\begin{aligned} -A(w) &= \int_{\mathbb{R}_+^n} \left( \langle dw, du_{(\xi, \epsilon)} \rangle_g + c_n R_g w u_{(\xi, \epsilon)} \right) + \int_{\partial \mathbb{R}_+^n} \left( d_n \kappa_g w u_{(\xi, \epsilon)} - nu_{(\xi, \epsilon)}^{\frac{n}{n-2}} w \right) \\ &= \int_{\mathbb{R}_+^n} f u_{(\xi, \epsilon)} + \int_{\partial \mathbb{R}_+^n} \bar{f} u_{(\xi, \epsilon)}, \end{aligned}$$

since  $u_{(\xi, \epsilon)} \in \Sigma_{(\xi, \epsilon)}$ . Then (2.12) implies

$$\begin{aligned} \frac{\theta}{2} \|w\|_\Sigma^2 &\leq \int_{\mathbb{R}_+^n} \left( |dw|_g^2 + c_n R_g w^2 \right) + \int_{\partial \mathbb{R}_+^n} \left( d_n \kappa_g w^2 - nu_{(\xi, \epsilon)}^{\frac{2}{n-2}} w^2 \right) + \frac{1}{\theta} A(w)^2 \\ &= \int_{\mathbb{R}_+^n} f w + \int_{\partial \mathbb{R}_+^n} \bar{f} w + \frac{1}{\theta} \left( \int_{\mathbb{R}_+^n} f u_{(\xi, \epsilon)} + \int_{\partial \mathbb{R}_+^n} \bar{f} u_{(\xi, \epsilon)} \right)^2 \\ &\leq \left\{ \|w\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)} + \frac{2}{\theta} \|u_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)}^2 \|f\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \right\} \|f\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \\ &\quad + \left\{ \|w\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)} + \frac{2}{\theta} \|u_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)}^2 \|\bar{f}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)} \right\} \|\bar{f}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)} \\ &\leq \left\{ K^{\frac{1}{2}} \|w\|_\Sigma + \frac{2}{\theta} \|u_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)}^2 \|f\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \right\} \|f\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \\ &\quad + \left\{ K^{\frac{1}{2}} \|w\|_\Sigma + \frac{2}{\theta} \|u_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)}^2 \|\bar{f}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)} \right\} \|\bar{f}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)}. \end{aligned}$$

Hence,

$$\begin{aligned} \frac{\theta}{4} \|w\|_\Sigma^2 &\leq \left\{ \frac{K}{\theta} + \frac{2}{\theta} \|u_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)}^2 \right\} \|f\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)}^2 \\ &\quad + \left\{ \frac{K}{\theta} + \frac{2}{\theta} \|u_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)}^2 \right\} \|\bar{f}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)}^2 \end{aligned}$$

and the result follows.  $\square$

**Proposition 2.8.** *Let  $\alpha_0$  be the constant obtained in Corollary 2.6. There is a constant  $\alpha_1 = \alpha_1(n)$ ,  $0 < \alpha_1 \leq \alpha_0$ , with the following property: if  $|h(x)| + |\partial h(x)| + |\partial^2 h(x)| \leq \alpha_1$  for all  $x \in \mathbb{R}_+^n$ , given any pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$  there exists a unique  $v_{(\xi, \epsilon)} \in \Sigma$  such that  $v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)} \in \Sigma_{(\xi, \epsilon)}$  and*

$$\int_{\mathbb{R}_+^n} \left( \langle dv_{(\xi, \epsilon)}, d\psi \rangle_g + c_n R_g v_{(\xi, \epsilon)} \psi \right) + \int_{\partial \mathbb{R}_+^n} \left( d_n \kappa_g v_{(\xi, \epsilon)} \psi - (n-2) |v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} v_{(\xi, \epsilon)} \psi \right) = 0$$

for all  $\psi \in \Sigma_{(\xi, \epsilon)}$ . Moreover, there exists  $C = C(n) > 0$  such that

$$\|v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}\|_\Sigma \leq C \|\Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} + C \|d_n \kappa_g u_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)} \quad (2.17)$$

In particular,  $v_{(\xi, \epsilon)} \neq 0$ .

*Proof.* Using Proposition 2.7 we can define

$$\mathcal{G}_{(\xi, \epsilon)} : L^{\frac{2n}{n+2}}(\mathbb{R}_+^n) \times L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n) \longrightarrow \Sigma_{(\xi, \epsilon)}$$

by  $\mathcal{G}_{(\xi, \epsilon)}(f, \bar{f}) = w$ , where  $w \in \Sigma_{(\xi, \epsilon)}$  satisfies (2.16) for all  $\psi \in \Sigma_{(\xi, \epsilon)}$ . Hence, there exists  $C = C(n)$  such that

$$\|\mathcal{G}_{(\xi, \epsilon)}(f, \bar{f})\|_\Sigma \leq C \|f\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} + C \|\bar{f}\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)}. \quad (2.18)$$

We define a nonlinear mapping  $\Phi_{(\xi, \epsilon)}(w) : \Sigma_{(\xi, \epsilon)} \rightarrow \Sigma_{(\xi, \epsilon)}$  by

$$\Phi_{(\xi, \epsilon)}(w) = \mathcal{G}_{(\xi, \epsilon)}(f_{(\xi, \epsilon)}, \bar{f}_{(\xi, \epsilon, w)})$$

where

$$f_{(\xi, \epsilon)} = \Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)}$$

and

$$\bar{f}_{(\xi, \epsilon, w)} = -d_n \kappa_g u_{(\xi, \epsilon)} + (n-2) \left\{ |u_{(\xi, \epsilon)} + w|^{\frac{2}{n-2}} (u_{(\xi, \epsilon)} + w) - u_{(\xi, \epsilon)}^{\frac{n}{n-2}} - \frac{n}{n-2} u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w \right\}.$$

It follows from Proposition 2.2 and the inequality (2.18) that  $\|\Phi_{(\xi, \epsilon)}(0)\|_\Sigma \leq C\alpha_1$ . Since

$$\begin{aligned} & \left| |u_{(\xi, \epsilon)} + w|^{\frac{2}{n-2}} (u_{(\xi, \epsilon)} + w) - |u_{(\xi, \epsilon)} + \tilde{w}|^{\frac{2}{n-2}} (u_{(\xi, \epsilon)} + \tilde{w}) - \frac{n}{n-2} u_{(\xi, \epsilon)}^{\frac{2}{n-2}} (w - \tilde{w}) \right| \\ & \leq C \left( |w|^{\frac{2}{n-2}} + |\tilde{w}|^{\frac{2}{n-2}} \right) |w - \tilde{w}|, \end{aligned}$$

we have

$$\begin{aligned} & \|\Phi_{(\xi, \epsilon)}(w) - \Phi_{(\xi, \epsilon)}(\tilde{w})\|_\Sigma \\ & \leq C \left\| \left( |w|^{\frac{2}{n-2}} + |\tilde{w}|^{\frac{2}{n-2}} \right) (w - \tilde{w}) \right\|_{L^{\frac{2(n-1)}{n}}(\partial \mathbb{R}_+^n)} \\ & \leq C \left\{ \|w\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)}^{\frac{2}{n-2}} + \|\tilde{w}\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)}^{\frac{2}{n-2}} \right\} \|w - \tilde{w}\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)} \end{aligned}$$

for all  $w, \tilde{w} \in \Sigma_{(\xi, \epsilon)}$ . Hence, it follows from the estimate (2.3) that

$$\|\Phi_{(\xi, \epsilon)}(w) - \Phi_{(\xi, \epsilon)}(\tilde{w})\|_{\Sigma} \leq C \left( \|w\|_{\Sigma}^{\frac{2}{n-2}} + \|\tilde{w}\|_{\Sigma}^{\frac{2}{n-2}} \right) \|w - \tilde{w}\|_{\Sigma}$$

for any  $w, \tilde{w} \in \Sigma_{(\xi, \epsilon)}$ . Thus, for  $\alpha_1$  small, the contraction maximum principle implies that the mapping  $\Phi_{(\xi, \epsilon)}$  has a fixed point  $w_{(\xi, \epsilon)}$ . Now the result follows from choosing  $v_{(\xi, \epsilon)} = u_{(\xi, \epsilon)} + w_{(\xi, \epsilon)}$ . Observe that  $v_{(\xi, \epsilon)}$  cannot be identically zero because of (2.17) and Proposition 2.2 with  $\alpha = \alpha_1$  small.  $\square$

Given a pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$  we define

$$\begin{aligned} \mathcal{F}_g(\xi, \epsilon) &= \int_{\mathbb{R}_+^n} (|dv_{(\xi, \epsilon)}|_g^2 + c_n R_g v_{(\xi, \epsilon)}^2) + \int_{\partial \mathbb{R}_+^n} d_n \kappa_g v_{(\xi, \epsilon)}^2 \\ &\quad - \frac{(n-2)^2}{n-1} \int_{\partial \mathbb{R}_+^n} |v_{(\xi, \epsilon)}|^{\frac{2(n-1)}{n-2}} - \frac{n-2}{n-1} \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{2(n-1)}{n-2}}. \end{aligned} \quad (2.19)$$

**Proposition 2.9.** *Suppose that  $|h(x)| + |\partial h(x)| + |\partial^2 h(x)| \leq \alpha_1$  for all  $x \in \mathbb{R}_+^n$ , where  $\alpha_1$  is the constant obtained in Proposition 2.8. Choosing  $\alpha_1$  smaller if necessary, the function  $\mathcal{F}_g$  is continuously differentiable and, if  $(\bar{\xi}, \bar{\epsilon})$  is a critical point of  $\mathcal{F}_g$ , then  $v_{(\bar{\xi}, \bar{\epsilon})}$  is a positive smooth solution of*

$$\begin{cases} \Delta_g v_{(\bar{\xi}, \bar{\epsilon})} - c_n R_g v_{(\bar{\xi}, \bar{\epsilon})} = 0, & \text{in } \mathbb{R}_+^n, \\ \frac{\partial}{\partial x_n} v_{(\bar{\xi}, \bar{\epsilon})} - d_n \kappa_g v_{(\bar{\xi}, \bar{\epsilon})} + (n-2)v_{(\bar{\xi}, \bar{\epsilon})}^{\frac{n}{n-2}} = 0, & \text{on } \partial \mathbb{R}_+^n. \end{cases} \quad (2.20)$$

In the proof of Proposition 2.9 we will use the following removable singularities theorem, which is a slight modification of Proposition 2.7 of [22]:

**Lemma 2.10.** *Let  $(M^n, g)$  be a Riemannian manifold with boundary  $\partial M$ . Let  $x \in \partial M$  be a boundary point and  $\mathcal{U} \subset M$  an open set containing  $x$ . Let  $u$  be a weak solution to*

$$\begin{cases} \Delta_g u + \phi u = 0, & \text{in } \mathcal{U} \setminus \{x\} \\ \frac{\partial u}{\partial \eta} + \psi u = 0, & \text{on } \mathcal{U} \cap \partial M \setminus \{x\}, \end{cases}$$

where  $\phi \in L^{\frac{n}{2}}(\mathcal{U})$  and  $\psi \in L^{n-1}(\mathcal{U} \cap \partial M)$ . Suppose that  $u \in L^q(\mathcal{U}) \cap L^p(\mathcal{U} \cap \partial M)$  for some  $q > \frac{n}{n-2}$  and  $p > \frac{n-1}{n-2}$ . Then  $u$  is a weak solution to

$$\begin{cases} \Delta_g u + \phi u = 0, & \text{in } \mathcal{U}, \\ \frac{\partial u}{\partial \eta} + \psi u = 0, & \text{on } \mathcal{U} \cap \partial M. \end{cases}$$

*Proof of Proposition 2.9.* Given a pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$ , by the definition of  $v_{(\xi, \epsilon)}$ , there exist  $b_a(\xi, \epsilon) \in \mathbb{R}$ ,  $a = 1, \dots, n$ , such that

$$\begin{aligned} &\int_{\mathbb{R}_+^n} (\langle dv_{(\xi, \epsilon)}, d\psi \rangle_g + c_n R_g v_{(\xi, \epsilon)} \psi) + \int_{\partial \mathbb{R}_+^n} (d_n \kappa_g v_{(\xi, \epsilon)} \psi - (n-2)|v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} v_{(\xi, \epsilon)} \psi) \\ &= \sum_{a=1}^n b_a(\xi, \epsilon) \cdot \int_{\partial \mathbb{R}_+^n} \phi_{(\xi, \epsilon, a)} \psi \end{aligned}$$

for any  $\psi \in \Sigma$ . Hence, derivating the expression (2.19) and observing the identity (2.2), we obtain

$$\frac{\partial \mathcal{F}_g}{\partial \epsilon}(\xi, \epsilon) = 2 \sum_{a=1}^n b_a(\xi, \epsilon) \cdot \int_{\partial \mathbb{R}_+^n} \phi_{(\xi, \epsilon, b)} \frac{\partial}{\partial \epsilon} v_{(\xi, \epsilon)}$$

and

$$\frac{\partial \mathcal{F}_g}{\partial \xi_k}(\xi, \epsilon) = 2 \sum_{a=1}^n b_a(\xi, \epsilon) \cdot \int_{\partial \mathbb{R}_+^n} \phi_{(\xi, \epsilon, a)} \frac{\partial}{\partial \xi_k} v_{(\xi, \epsilon)}, \quad k = 1, \dots, n-1.$$

On the other hand,

$$\int_{\partial \mathbb{R}_+^n} \phi_{(\xi, \epsilon, a)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) = 0, \quad a = 1, \dots, n,$$

since  $v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)} \in \Sigma_{(\xi, \epsilon)}$ . This implies

$$\begin{aligned} 0 &= \int_{\partial \mathbb{R}_+^n} \frac{\partial}{\partial \epsilon} \phi_{(\xi, \epsilon, a)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) + \int_{\partial \mathbb{R}_+^n} \phi_{(\xi, \epsilon, a)} \frac{\partial}{\partial \epsilon} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \\ &= \int_{\partial \mathbb{R}_+^n} \frac{\partial}{\partial \epsilon} \phi_{(\xi, \epsilon, a)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) + \int_{\partial \mathbb{R}_+^n} \phi_{(\xi, \epsilon, a)} \frac{\partial}{\partial \epsilon} v_{(\xi, \epsilon)} + \beta(n) \delta_{an} \epsilon^{-1} \end{aligned}$$

and

$$0 = \int_{\partial \mathbb{R}_+^n} \frac{\partial}{\partial \xi_k} \phi_{(\xi, \epsilon, a)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) + \int_{\partial \mathbb{R}_+^n} \phi_{(\xi, \epsilon, a)} \frac{\partial}{\partial \xi_k} v_{(\xi, \epsilon)} - \beta(n) \delta_{ak} \epsilon^{-1},$$

where

$$\beta(n) = -\epsilon \int_{\partial \mathbb{R}_+^n} \phi_{(\xi, \epsilon, n)} \frac{\partial}{\partial \epsilon} u_{(\xi, \epsilon)} = \epsilon \int_{\partial \mathbb{R}_+^n} \phi_{(\xi, \epsilon, k)} \frac{\partial}{\partial \xi_k} u_{(\xi, \epsilon)} > 0, \quad k = 1, \dots, n-1.$$

Thus

$$-b_n(\xi, \epsilon) \beta(n) = \frac{\epsilon}{2} \frac{\partial \mathcal{F}_g}{\partial \epsilon}(\xi, \epsilon) + \epsilon \sum_{a=1}^n b_a(\xi, \epsilon) \cdot \int_{\partial \mathbb{R}_+^n} \frac{\partial}{\partial \epsilon} \phi_{(\xi, \epsilon, a)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}).$$

Similarly,

$$b_k(\xi, \epsilon) \beta(n) = \frac{\epsilon}{2} \frac{\partial \mathcal{F}_g}{\partial \xi_k}(\xi, \epsilon) + \epsilon \sum_{a=1}^n b_a(\xi, \epsilon) \cdot \int_{\partial \mathbb{R}_+^n} \frac{\partial}{\partial \xi_k} \phi_{(\xi, \epsilon, a)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)})$$

for  $k = 1, \dots, n-1$ . Hence, if  $(\bar{\xi}, \bar{\epsilon})$  is a critical point of  $\mathcal{F}_g$ , then there exists  $C = C(n)$  such that

$$\sum_{a=1}^n |b_a(\bar{\xi}, \bar{\epsilon})| \leq C \|v_{(\bar{\xi}, \bar{\epsilon})} - u_{(\bar{\xi}, \bar{\epsilon})}\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)} \sum_{a=1}^n |b_a(\bar{\xi}, \bar{\epsilon})|.$$

By the estimate (2.3) and Propositions 2.2 and 2.8,  $\|v_{(\bar{\xi}, \bar{\varepsilon})} - u_{(\bar{\xi}, \bar{\varepsilon})}\|_{L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n)} \leq CK^{\frac{1}{2}}\alpha_1$ . Thus, choosing  $\alpha_1$  small, we must have  $b_a(\bar{\xi}, \bar{\varepsilon}) = 0$  for  $a = 1, \dots, n$ . Hence,

$$\begin{aligned} & \int_{\mathbb{R}_+^n} \left( \langle dv_{(\bar{\xi}, \bar{\varepsilon})}, d\psi \rangle_g + c_n R_g v_{(\bar{\xi}, \bar{\varepsilon})} \psi \right) \\ & + \int_{\partial\mathbb{R}_+^n} \left( d_n \kappa_g v_{(\bar{\xi}, \bar{\varepsilon})} \psi - (n-2) |v_{(\bar{\xi}, \bar{\varepsilon})}|^{\frac{2}{n-2}} v_{(\bar{\xi}, \bar{\varepsilon})} \psi \right) = 0 \end{aligned} \quad (2.21)$$

for any  $\psi \in \Sigma$ .

Now we are going to show that  $v_{(\bar{\xi}, \bar{\varepsilon})} \geq 0$  on  $\partial\mathbb{R}_+^n$ . To that end, we set  $\psi = \min\{v_{(\bar{\xi}, \bar{\varepsilon})}, 0\}$  and use the equation (2.21) to conclude that

$$\begin{aligned} & \int_{\mathbb{R}_+^n \cap \{v_{(\bar{\xi}, \bar{\varepsilon})} < 0\}} \left( |dv_{(\bar{\xi}, \bar{\varepsilon})}|_g^2 + c_n R_g v_{(\bar{\xi}, \bar{\varepsilon})}^2 \right) \\ & + \int_{\partial\mathbb{R}_+^n \cap \{v_{(\bar{\xi}, \bar{\varepsilon})} < 0\}} d_n \kappa_g v_{(\bar{\xi}, \bar{\varepsilon})}^2 = (n-2) \int_{\partial\mathbb{R}_+^n \cap \{v_{(\bar{\xi}, \bar{\varepsilon})} < 0\}} |v_{(\bar{\xi}, \bar{\varepsilon})}|^{\frac{2(n-1)}{n-2}}. \end{aligned} \quad (2.22)$$

Using (2.11) with  $w = \psi$  we see that

$$\begin{aligned} \left( \int_{\partial\mathbb{R}_+^n \cap \{v_{(\bar{\xi}, \bar{\varepsilon})} < 0\}} |v_{(\bar{\xi}, \bar{\varepsilon})}|^{\frac{2(n-1)}{n-2}} \right)^{\frac{n-2}{n-1}} & \leq 2K \int_{\mathbb{R}_+^n \cap \{v_{(\bar{\xi}, \bar{\varepsilon})} < 0\}} \left( |dv_{(\bar{\xi}, \bar{\varepsilon})}|_g^2 + c_n R_g v_{(\bar{\xi}, \bar{\varepsilon})}^2 \right) \\ & + 2K \int_{\partial\mathbb{R}_+^n \cap \{v_{(\bar{\xi}, \bar{\varepsilon})} < 0\}} d_n \kappa_g v_{(\bar{\xi}, \bar{\varepsilon})}^2. \end{aligned}$$

From this, together with (2.22), we deduce that  $v_{(\bar{\xi}, \bar{\varepsilon})} \geq 0$  almost everywhere on  $\partial\mathbb{R}_+^n$  or

$$\left( \int_{\partial\mathbb{R}_+^n \cap \{v_{(\bar{\xi}, \bar{\varepsilon})} < 0\}} |v_{(\bar{\xi}, \bar{\varepsilon})}|^{\frac{2(n-1)}{n-2}} \right)^{\frac{1}{n-1}} \geq \frac{1}{2K(n-2)}.$$

On the other hand,

$$\left( \int_{\partial\mathbb{R}_+^n \cap \{v_{(\bar{\xi}, \bar{\varepsilon})} < 0\}} |v_{(\bar{\xi}, \bar{\varepsilon})}|^{\frac{2(n-1)}{n-2}} \right)^{\frac{n-2}{2(n-1)}} \leq \left( \int_{\partial\mathbb{R}_+^n} |v_{(\bar{\xi}, \bar{\varepsilon})} - u_{(\bar{\xi}, \bar{\varepsilon})}|^{\frac{2(n-1)}{n-2}} \right)^{\frac{n-2}{2(n-1)}} \leq CK^{\frac{1}{2}}\alpha_1.$$

Hence, choosing  $\alpha_1$  sufficiently small we have  $v_{(\bar{\xi}, \bar{\varepsilon})} \geq 0$  on  $\partial\mathbb{R}_+^n$ . In particular, the equation (2.21) can be written as

$$\int_{\mathbb{R}_+^n} \left( \langle dv_{(\bar{\xi}, \bar{\varepsilon})}, d\psi \rangle_g + c_n R_g v_{(\bar{\xi}, \bar{\varepsilon})} \psi \right) + \int_{\partial\mathbb{R}_+^n} \left( d_n \kappa_g v_{(\bar{\xi}, \bar{\varepsilon})} \psi - (n-2) |v_{(\bar{\xi}, \bar{\varepsilon})}|^{\frac{n}{n-2}} \psi \right) = 0$$

for any  $\psi \in \Sigma$ . By a result of Cherrier in [10],  $v_{(\bar{\xi}, \bar{\varepsilon})}$  is smooth.

The fact that  $v_{(\bar{\xi}, \bar{\epsilon})} > 0$  in  $\mathbb{R}_+^n$  is just a consequence of the maximum principle, as follows. We set  $\tilde{g} = \tilde{u}^{-\frac{4}{n-2}} g$ , where  $\tilde{u}(x) = (1 + |x|^2)^{\frac{2-n}{2}}$ . Observe that  $\tilde{u}$  satisfies  $\Delta \tilde{u} + n(n-2)\tilde{u}^{-\frac{n+2}{n-2}} = 0$  in  $\mathbb{R}_+^n$  and we have

$$\begin{aligned} c_n R_{\tilde{g}} &= -\tilde{u}^{-\frac{n+2}{n-2}} \Delta \tilde{u} - \tilde{u}^{-\frac{n+2}{n-2}} (\Delta_g \tilde{u} - \Delta \tilde{u} - c_n R_g \tilde{u}) \\ &\geq n(n-2) - C \tilde{u}^{-\frac{n+2}{n-2}} \left\{ |h| |\partial^2 \tilde{u}| + |\partial h| |\partial \tilde{u}| + (|\partial^2 h| + |\partial h|^2) |\tilde{u}| \right\}. \end{aligned}$$

Using the facts that  $h(x) = 0$  for  $|x| \geq 1$  and  $|h| + |\partial h| + |\partial^2 h| \leq C\alpha_1$  we can assume that  $R_{\tilde{g}} > 0$ , by choosing  $\alpha_1$  small.

Let  $S_+^n$  be a hemisphere of  $S_{1/2}^n$ . We will use the well known conformal equivalence between  $S_+^n \setminus \{x_0\}$  and  $\mathbb{R}_+^n$  realized by the stereographic projection, where  $x_0 \in \partial S_+^n$ . Under this equivalence, the standard metric on  $S_+^n$  is written on  $\mathbb{R}_+^n$  as  $\tilde{u}^{-\frac{4}{n-2}} \delta$ , where  $\delta$  is the Euclidean metric on  $\mathbb{R}_+^n$ . We set  $\tilde{v} = \tilde{u}^{-1} v_{(\bar{\xi}, \bar{\epsilon})}$ . By the properties (2.8) of the operators  $L_g = \Delta_g - c_n R_g$  and  $B_g = \frac{\partial}{\partial \eta} - d_n \kappa_g$ , we have

$$L_{\tilde{g}}(\tilde{v}) = \tilde{u}^{-\frac{n+2}{n-2}} L_g v_{(\bar{\xi}, \bar{\epsilon})} = 0, \quad \text{in } S_+^n,$$

and

$$B_{\tilde{g}}(\tilde{v}) + (n-2)\tilde{v}^{-\frac{n}{n-2}} = \tilde{u}^{-\frac{n}{n-2}} B_g v_{(\bar{\xi}, \bar{\epsilon})} + (n-2)(\tilde{u}^{-1} v_{(\bar{\xi}, \bar{\epsilon})})^{-\frac{n}{n-2}} = 0, \quad \text{on } \partial S_+^n.$$

To establish the last two equations, we also used Lemma 2.10.

Since  $R_{\tilde{g}} > 0$ , it follows from the maximum principle in  $S_+^n$  and the Hopf Lemma that if  $\tilde{v} \geq 0$  on  $\partial S_+^n$  then we have either  $\tilde{v} > 0$  or  $\tilde{v} \equiv 0$  in  $S_+^n$ . The latter contradicts the last assertion of Proposition 2.8. Hence,  $\tilde{v} \geq 0$  on  $\partial S_+^n$  implies that  $\tilde{v} > 0$  in  $S_+^n$ . Since we have proved that  $v_{(\bar{\xi}, \bar{\epsilon})} \geq 0$  on  $\partial \mathbb{R}_+^n$ , we conclude that  $v_{(\bar{\xi}, \bar{\epsilon})} > 0$  in  $\mathbb{R}_+^n$ .  $\square$

### 3 An estimate for the energy of a bubble

In this section we will show that the energy function  $\mathcal{F}_g$  can be approximated by a certain auxiliary function.

We fix a multi-linear form  $W : \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  satisfying the algebraic properties of the Weyl tensor. We set

$$|W|^2 = \sum_{a,b,c,d=1}^n (W_{abcd} + W_{adbc})^2$$

and assume that  $|W|^2 > 0$ . Recall that throughout this article we work with indices  $1 \leq i, j, k, l \leq n-1$  and  $1 \leq a, b, c, d \leq n$  and set  $\bar{x} = (x_1, \dots, x_{n-1}, 0) \in \partial \mathbb{R}_+^n$  whenever  $x = (x_1, \dots, x_{n-1}, x_n) \in \mathbb{R}_+^n$ . For  $x \in \mathbb{R}_+^n$  we set

$$H_{ij}(x) = H_{ij}(\bar{x}) = W_{ikjl} x^k x^l \quad \text{and} \quad H_{nb}(x) = 0$$



and define  $\bar{H}_{ab}(x) = f(|\bar{x}|^2)H_{ab}(x)$ , where

$$f(s) = \sum_{j=0}^d a_j s^j. \quad (3.1)$$

The integer  $0 < d < \frac{n-6}{4}$  and the coefficients  $a_0, \dots, a_d \in \mathbb{R}$  will be chosen later. Observe that  $H$  is symmetric, trace-free, independent of the coordinate  $x_n$  and satisfies

$$x^a H_{ab}(x) = x^i H_{ib}(x) = 0 = \partial_a H_{ab}(x) = \partial_i H_{ib}(x), \quad \text{for any } x \in \mathbb{R}_+^n.$$

We define a Riemannian metric  $g = \exp(h)$  on  $\mathbb{R}_+^n$  where  $h$  is a trace-free symmetric two tensor on  $\mathbb{R}_+^n$  satisfying

$$\begin{cases} h_{ab}(x) = \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ab}(x), & \text{for } |x| \leq \rho, \\ h_{ab}(x) = 0, & \text{for } |x| \geq 1. \end{cases}$$

Here,  $\mu \leq 1$ ,  $\lambda \leq \rho \leq 1$  and we suppose that  $h_{nb}(x) = 0$  for any  $x \in \mathbb{R}_+^n$  and  $\partial_n h_{ab}(x) = 0$  for any  $x \in \partial \mathbb{R}_+^n$ . We also assume that  $|h| + |\partial h| + |\partial^2 h| \leq \alpha_1$  where  $\alpha_1$  is the constant obtained in Proposition 2.8. Observe that

$$x^a h_{ab}(x) = x^i h_{ib}(x) = 0 = \partial_a h_{ab}(x) = \partial_i h_{ib}(x), \quad \text{for } |x| \leq \rho.$$

and  $h_{ab}(x) = O(\mu(\lambda + |x|)^{2d+2})$ . The second fundamental form of  $\partial \mathbb{R}_+^n$  satisfies

$$\pi_{ij} = \Gamma_{ij}^n = \frac{1}{2}(g_{m,j} + g_{j,n,i} - g_{ij,n}) = 0.$$

In particular, the mean curvature of  $\partial \mathbb{R}_+^n$  is given by  $\kappa_g = \frac{1}{n-1} g^{ij} \pi_{ij} = 0$ .

Using Proposition 2.8, for each pair  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$  we choose  $v_{(\xi, \epsilon)}$  to be the unique element of  $\Sigma$  such that  $v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)} \in \Sigma_{(\xi, \epsilon)}$  and

$$\int_{\mathbb{R}_+^n} (\langle dv_{(\xi, \epsilon)}, d\psi \rangle_g + c_n R_g v_{(\xi, \epsilon)} \psi) - (n-2) \int_{\partial \mathbb{R}_+^n} |v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} v_{(\xi, \epsilon)} \psi = 0$$

for all  $\psi \in \Sigma_{(\xi, \epsilon)}$ .

Finally, we define  $\Omega = \{(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty); |\xi| < 1, \frac{1}{2} < \epsilon < 2\}$ .

**Proposition 3.1.** *For any pair  $(\xi, \epsilon) \in \lambda \Omega$  we have the estimates*

$$\left\| \Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)} \right\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \leq C \mu \lambda^{2d+2} + C \left( \frac{\lambda}{\rho} \right)^{\frac{n-2}{2}}$$

and

$$\left\| \Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)} + \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} \right\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \leq C \mu^2 \lambda^{4d+4} + C \left( \frac{\lambda}{\rho} \right)^{\frac{n-2}{2}}.$$

*Proof.* We just observe that

$$|\Delta_g u_{(\xi, \epsilon)}(x) - c_n R_g(x) u_{(\xi, \epsilon)}(x)| \leq C \mu \lambda^{\frac{n-2}{2}} (\lambda + |x|)^{2d+2-n}$$

and

$$|\Delta_g u_{(\xi, \epsilon)}(x) - c_n R_g(x) u_{(\xi, \epsilon)}(x) + \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ij}(x) \partial_i \partial_j u_{(\xi, \epsilon)}(x)| \leq C \mu^2 \lambda^{\frac{n-2}{2}} (\lambda + |x|)^{4d+4-n}$$

for  $|x| \leq \rho$ . In the last inequality we used the fact that, since  $\partial_a h_{ab}(x) = 0$  for  $|x| \leq \rho$ , Lemma 2.1 implies that  $|R_g(x)| \leq |\partial h(x)|^2 + |h(x)| |\partial^2 h(x)|$  for  $|x| \leq \rho$ .  $\square$

**Corollary 3.2.** *For any pair  $(\xi, \epsilon) \in \lambda\Omega$  we have the estimate*

$$\|v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)} + \|v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n)} \leq C \mu \lambda^{2d+2} + C \left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}}.$$

*Proof.* It follows from Proposition 2.8 and the estimate (2.3) that

$$\begin{aligned} \|v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)} + \|v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n)} &\leq C \|\Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \\ &\leq C \mu \lambda^{2d+2} + C \left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}}, \end{aligned}$$

where we used Proposition 3.1 in the last inequality.  $\square$

In order to refine the estimate of Corollary 3.2, using Proposition 2.7 with  $h_{ab} = 0$  we choose the function  $w_{(\xi, \epsilon)}$  to be the unique element of  $\Sigma_{(\xi, \epsilon)}$  satisfying

$$\int_{\mathbb{R}_+^n} \langle dw_{(\xi, \epsilon)}, d\psi \rangle - \int_{\partial\mathbb{R}_+^n} n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w_{(\xi, \epsilon)} \psi = - \int_{\mathbb{R}_+^n} \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} \psi \quad (3.2)$$

for all  $\psi \in \Sigma_{(\xi, \epsilon)}$ . Observe that, since  $x^i H_{ij}(x) = 0$  for any  $x \in \mathbb{R}_+^n$ , we have  $w_{(0, \epsilon)} = 0$ .

**Proposition 3.3.** *The function  $w_{(\xi, \epsilon)}$  is smooth and satisfies, for any pair  $(\xi, \epsilon) \in \lambda\Omega$ ,*

$$|\partial^k w_{(\xi, \epsilon)}(x)| \leq C \lambda^{\frac{n-2}{2}} \mu (\lambda + |x|)^{2d+4-k-n}, \quad \text{for all } x \in \mathbb{R}_+^n, \quad k = 0, 1, 2.$$

*Proof.* First observe that there exist real numbers  $b_a(\xi, \epsilon)$ ,  $1 \leq a \leq n$ , such that  $w_{(\xi, \epsilon)}$  satisfies

$$\begin{aligned} \int_{\mathbb{R}_+^n} \langle dw_{(\xi, \epsilon)}, d\psi \rangle - \int_{\partial\mathbb{R}_+^n} n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w_{(\xi, \epsilon)} \psi & \\ = - \int_{\mathbb{R}_+^n} \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} \psi + \sum_{a=1}^n b_a(\xi, \epsilon) \int_{\partial\mathbb{R}_+^n} \phi_{(a, \xi, \epsilon)} \psi & \end{aligned} \quad (3.3)$$

for all  $\psi \in \Sigma$ . Hence, it follows from standard elliptic theory that  $w_{(\xi, \epsilon)}$  is smooth.

Now we are going to prove the pointwise estimates. Observe that

$$\left\| \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ij}(x) \partial_i \partial_j u_{(\xi, \epsilon)}(x) \right\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \leq C \mu \lambda^{2d+2}. \quad (3.4)$$

Then we apply Proposition 2.7 with  $h_{ab} = 0$  and use the estimates (2.3) and (3.4) to conclude that

$$\|w_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)} + \|w_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n)} \leq K^{\frac{1}{2}} \|w_{(\xi, \epsilon)}\|_{\Sigma} \leq C \mu \lambda^{2d+2}.$$

Moreover, we can use the equation (3.3) with  $\psi = \phi_{(\xi, \epsilon, \rho)}$  to conclude that

$$\sum_{a=0}^n |b_a(\xi, \epsilon)| \leq C \mu \lambda^{2d+2}.$$

Hence,

$$|\Delta w_{(\xi, \epsilon)}(x)| = \left| \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ij}(x) \partial_i \partial_j u_{(\xi, \epsilon)}(x) \right| \leq \mu \lambda^{\frac{n-2}{2}} (\lambda + |x|)^{2d+2-n},$$

for all  $x \in \mathbb{R}_+^n$ , and

$$\left| \frac{\partial}{\partial x_n} w_{(\xi, \epsilon)}(x) + n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w_{(\xi, \epsilon)}(x) \right| = \left| - \sum_{a=1}^n b_a(\xi, \epsilon) \phi_{(a, \xi, \epsilon)}(x) \right| \leq \mu \lambda^{\frac{n}{2}} (\lambda + |x|)^{2d+2-n}$$

for all  $x \in \partial\mathbb{R}_+^n$ .

*Claim.*  $\sup_{x \in \mathbb{R}_+^n} (\lambda + |x|)^{\frac{n-2}{2}} |w_{(\xi, \epsilon)}(x)| \leq C \mu \lambda^{2d+2}$

We fix  $x_0 \in \mathbb{R}_+^n$  and set  $r = \frac{1}{2}(\lambda + |x_0|)$ . Then we see that

$$u_{(\xi, \epsilon)}^{\frac{2}{n-2}}(x) \leq C r^{-1}, \quad \text{for all } x \in B_r^+(x_0),$$

$$\left| \frac{\partial}{\partial x_n} w_{(\xi, \epsilon)}(x) + n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w_{(\xi, \epsilon)}(x) \right| \leq C \mu \lambda^{\frac{n}{2}} r^{2d+2-n}, \quad \text{for all } x \in B_r^+(x_0) \cap \partial\mathbb{R}_+^n$$

and

$$|\Delta w_{(\xi, \epsilon)}(x)| \leq C \mu \lambda^{\frac{n-2}{2}} r^{2d+2-n}, \quad \text{for all } x \in B_r^+(x_0).$$

It follows from standard interior estimates that

$$\begin{aligned} r^{\frac{n-2}{2}} |w_{(\xi, \epsilon)}(x_0)| &\leq C \|w_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n-2}}(B_r^+(x_0))} + C r^{\frac{n+2}{2}} \|\Delta w_{(\xi, \epsilon)}\|_{L^\infty(B_r^+(x_0))} \\ &\quad + C r^{\frac{n}{2}} \left\| \frac{\partial}{\partial x_n} w_{(\xi, \epsilon)} + n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} w_{(\xi, \epsilon)} \right\|_{L^\infty(B_r^+(x_0) \cap \partial\mathbb{R}_+^n)} \\ &\leq C \mu \lambda^{2d+2} + C \mu \lambda^{\frac{n-2}{2}} r^{2d+2+\frac{2-n}{2}} + C \mu \lambda^{\frac{n}{2}} r^{2d+2-\frac{n}{2}} \\ &\leq C \mu \lambda^{2d+2}, \end{aligned}$$

since we are assuming that  $d < \frac{n-6}{4}$ . This proves the Claim.

Since  $\sup_{x \in \mathbb{R}_+^n} |x|^{\frac{n-2}{2}} |w_{(\xi, \epsilon)}(x)| < \infty$ , for all  $x = (x_1, \dots, x_{n-1}, x_n) \in \mathbb{R}_+^n$  we have

$$\begin{aligned} w_{(\xi, \epsilon)}(x) &= -\frac{1}{(n-2)\sigma_{n-2}} \int_{\mathbb{R}_+^n} (|x-y|^{2-n} + |\tilde{x}-y|^{2-n}) \Delta w_{(\xi, \epsilon)}(y) dy \\ &\quad - \frac{1}{(n-2)\sigma_{n-2}} \int_{\partial \mathbb{R}_+^n} (|x-y|^{2-n} + |\tilde{x}-y|^{2-n}) \frac{\partial}{\partial y_n} w_{(\xi, \epsilon)}(y) dy, \end{aligned}$$

where  $\tilde{x} = (x_1, \dots, x_{n-1}, -x_n)$ . Now we use a bootstrap argument to prove the pointwise estimates. It follows from the last two inequalities that

$$\begin{aligned} \sup_{x \in \mathbb{R}_+^n} (\lambda + |x|)^\beta |w_{(\xi, \epsilon)}(x)| &\leq C \sup_{x \in \mathbb{R}_+^n} (\lambda + |x|^{\beta+2}) |\Delta w_{(\xi, \epsilon)}(x)| \\ &\quad + C \sup_{x \in \partial \mathbb{R}_+^n} (\lambda + |x|)^{\beta+1} \left| \frac{\partial}{\partial x_n} w_{(\xi, \epsilon)}(x) \right| \end{aligned}$$

for all  $0 < \beta < n-2$ . Since

$$|\Delta w_{(\xi, \epsilon)}(x)| \leq \mu \lambda^{\frac{n-2}{2}} (\lambda + |x|)^{2d+2-n}, \quad \text{for all } x \in \mathbb{R}_+^n,$$

and

$$\left| \frac{\partial}{\partial x_n} w_{(\xi, \epsilon)}(x) \right| \leq n \mu^{\frac{2}{n-2}} (x) |w_{(\xi, \epsilon)}(x)| + \mu \lambda^{\frac{n}{2}} (\lambda + |x|)^{2d+2-n}, \quad \text{for all } x \in \partial \mathbb{R}_+^n,$$

we see that

$$\sup_{x \in \mathbb{R}_+^n} (\lambda + |x|)^\beta |w_{(\xi, \epsilon)}(x)| \leq C \lambda \sup_{x \in \partial \mathbb{R}_+^n} (\lambda + |x|)^{\beta-1} |w_{(\xi, \epsilon)}(x)| + C \mu \lambda^{\beta+2d+3-\frac{n}{2}}$$

for all  $0 < \beta \leq n-4-2d$ . Interating we obtain

$$\sup_{x \in \mathbb{R}_+^n} (\lambda + |x|)^{n-2d-4} |w_{(\xi, \epsilon)}(x)| \leq C \mu \lambda^{\frac{n-2}{2}}.$$

The derivative estimates follow from elliptic theory, finishing the proof.  $\square$

**Corollary 3.4.** *For any  $(\xi, \epsilon) \in \lambda\Omega$ , the function  $v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)} - w_{(\xi, \epsilon)}$  satisfies*

$$\begin{aligned} &\|v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)} - w_{(\xi, \epsilon)}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)} + \|v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)} - w_{(\xi, \epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial \mathbb{R}_+^n)} \\ &\leq C \mu^{\frac{n}{n-2}} \lambda^{\frac{(2d+2)n}{n-2}} + C \left( \frac{\lambda}{\rho} \right)^{\frac{n-2}{2}}. \end{aligned}$$

*Proof.* It follows from the definition of  $w_{(\xi, \epsilon)}$  that

$$\begin{aligned} &\int_{\mathbb{R}_+^n} (\langle dw_{(\xi, \epsilon)}, d\psi \rangle_g + c_n R_g w_{(\xi, \epsilon)} \psi) - \int_{\partial \mathbb{R}_+^n} n \mu^{\frac{2}{n-2}} w_{(\xi, \epsilon)} \psi \\ &= - \int_{\mathbb{R}_+^n} \left\{ \partial_j \left( (g^{ij} - \delta_{ij}) \partial_i w_{(\xi, \epsilon)} \right) \psi - c_n R_g w_{(\xi, \epsilon)} \psi \right\} \\ &\quad - \int_{\mathbb{R}_+^n} \mu \lambda^{2d} f(\lambda^{-2} |\tilde{x}|^2) H_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} \psi, \end{aligned}$$

for any  $\psi \in \Sigma_{(\xi, \epsilon)}$ . Hence we can write  $w_{(\xi, \epsilon)} = -\mathcal{G}_{(\xi, \epsilon)}(B_1 + B_2, 0)$ , where

$$\begin{aligned} B_1 &= \partial_j \left( (g^{ij} - \delta_{ij}) \partial_i w_{(\xi, \epsilon)} \right) - c_n R_g w_{(\xi, \epsilon)}, \\ B_2 &= \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} \end{aligned}$$

and  $\mathcal{G}_{(\xi, \epsilon)}$  is the operator defined in the proof of Proposition 2.8.

On the other hand,

$$\begin{aligned} & \int_{\mathbb{R}_+^n} \left\{ \langle d(v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}), d\psi \rangle_g + c_n R_g (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \psi \right\} \\ & \quad - \int_{\partial \mathbb{R}_+^n} n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \psi \\ &= - \int_{\mathbb{R}_+^n} \left\{ \langle du_{(\xi, \epsilon)}, d\psi \rangle_g + c_n R_g u_{(\xi, \epsilon)} \psi \right\} \\ & \quad + \int_{\partial \mathbb{R}_+^n} \left\{ (n-2) |v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} v_{(\xi, \epsilon)} \psi - n u_{(\xi, \epsilon)}^{\frac{2}{n-2}} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \psi \right\} \\ &= \int_{\mathbb{R}_+^n} (\Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)}) \psi \\ & \quad + (n-2) \int_{\partial \mathbb{R}_+^n} \left\{ |v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}^{\frac{n}{n-2}} - \frac{n}{n-2} u_{(\xi, \epsilon)}^{\frac{2}{n-2}} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \right\} \psi. \end{aligned}$$

Hence we can write  $v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)} = \mathcal{G}_{(\xi, \epsilon)}(B_3, (n-2)B_4)$ , where

$$\begin{aligned} B_3 &= \Delta_g u_{(\xi, \epsilon)} - c_n R_g u_{(\xi, \epsilon)}, \\ B_4 &= (|v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}^{\frac{n}{n-2}}) - \frac{n}{n-2} u_{(\xi, \epsilon)}^{\frac{2}{n-2}} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}). \end{aligned}$$

Putting this facts together we conclude that

$$v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)} - w_{(\xi, \epsilon)} = \mathcal{G}_{(\xi, \epsilon)}(B_1 + B_2 + B_3, (n-2)B_4).$$

Now we are going to estimate the terms  $B_1, B_2, B_3, B_4$ . Since

$$\begin{aligned} |B_1(x)| &\leq C \partial(|h| |\partial w_{(\xi, \epsilon)}|)(x) + C(|\partial^2 h| |h| + |\partial h|^2) |w_{(\xi, \epsilon)}|(x) \\ &\leq C \mu^2 \lambda^{\frac{n-2}{2}} (\lambda + |x|)^{4d+4-n}, \quad \text{for } |x| \leq \rho, \end{aligned}$$

we have

$$\|B_1\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \leq C \mu^2 \lambda^{4d+4} + C \left( \frac{\lambda}{\rho} \right)^{\frac{n-2}{2}}.$$

It follows from Proposition 3.1 that

$$\|B_2 + B_3\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \leq C \mu^2 \lambda^{4d+4} + C \left( \frac{\lambda}{\rho} \right)^{\frac{n-2}{2}}.$$

Since  $|B_4(x)| \leq C|v_{(\xi,\epsilon)}(x) - u_{(\xi,\epsilon)}(x)|^{\frac{n}{n-2}}$  for any  $x \in \partial\mathbb{R}_+^n$ , we have

$$\|B_4\|_{L^{\frac{2(n-1)}{n}}(\partial\mathbb{R}_+^n)} \leq C\|v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n)}^{\frac{n}{n-2}} \leq C\mu^{\frac{n}{n-2}}\lambda^{\frac{(2d+2)n}{n-2}} + C\left(\frac{\lambda}{\rho}\right)^{\frac{n}{2}},$$

where in the last inequality we used Corollary 3.2.

Using the estimates above we see that

$$\|B_1 + B_2 + B_3\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} + \|(n-2)B_4\|_{L^{\frac{2(n-1)}{n}}(\partial\mathbb{R}_+^n)} \leq C\mu^{\frac{n}{n-2}}\lambda^{\frac{(2d+2)n}{n-2}} + C\left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}}.$$

Hence,

$$\|v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)} - w_{(\xi,\epsilon)}\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} + \|v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)} - w_{(\xi,\epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n)} \leq C\mu^{\frac{n}{n-2}}\lambda^{\frac{(2d+2)n}{n-2}} + C\left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}}.$$

□

**Lemma 3.5.** *For any  $(\xi, \epsilon) \in \lambda\Omega$  we have the estimate*

$$\left| \int_{\partial\mathbb{R}_+^n} (|v_{(\xi,\epsilon)}|^{\frac{2}{n-2}} - u_{(\xi,\epsilon)}^{\frac{2}{n-2}}) u_{(\xi,\epsilon)} v_{(\xi,\epsilon)} - \frac{1}{n-1} \int_{\partial\mathbb{R}_+^n} (|v_{(\xi,\epsilon)}|^{\frac{2(n-1)}{n-2}} - u_{(\xi,\epsilon)}^{\frac{2(n-1)}{n-2}}) \right| \leq C\mu^{\frac{2(n-1)}{n-2}} \lambda^{\frac{(4d+4)(n-1)}{n-2}} + C\left(\frac{\lambda}{\rho}\right)^{n-1}.$$

*Proof.* It follows from the pointwise estimate

$$\left| \left( |v_{(\xi,\epsilon)}|^{\frac{2}{n-2}} - u_{(\xi,\epsilon)}^{\frac{2}{n-2}} \right) u_{(\xi,\epsilon)} v_{(\xi,\epsilon)} - \frac{1}{n-1} \left( |v_{(\xi,\epsilon)}|^{\frac{2(n-1)}{n-2}} - u_{(\xi,\epsilon)}^{\frac{2(n-1)}{n-2}} \right) \right| \leq C|v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)}|^{\frac{2(n-1)}{n-2}}$$

that

$$\begin{aligned} & \left| \int_{\partial\mathbb{R}_+^n} (|v_{(\xi,\epsilon)}|^{\frac{2}{n-2}} - u_{(\xi,\epsilon)}^{\frac{2}{n-2}}) u_{(\xi,\epsilon)} v_{(\xi,\epsilon)} - \frac{1}{n-1} \int_{\partial\mathbb{R}_+^n} (|v_{(\xi,\epsilon)}|^{\frac{2(n-1)}{n-2}} - u_{(\xi,\epsilon)}^{\frac{2(n-1)}{n-2}}) \right| \\ & \leq C\|v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)}\|_{L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n)}^{\frac{2(n-1)}{n-2}} \leq C\left( \mu\lambda^{2d+2} + \left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}} \right)^{\frac{2(n-1)}{n-2}}, \end{aligned}$$

where in the last inequality we used Corollary 3.2. Now the result follows. □

**Proposition 3.6.** *Let  $\mathcal{F}_g$  be the function defined by the formula (2.19). For any pair  $(\xi, \epsilon) \in \lambda\Omega$  we have the estimate*

$$\begin{aligned} & \left| \mathcal{F}_g(\xi, \epsilon) - \frac{1}{2} \int_{B_\rho^+(0)} h_{ij} h_{ji} \partial_i u_{(\xi,\epsilon)} \partial_j u_{(\xi,\epsilon)} + \frac{c_n}{4} \int_{B_\rho^+(0)} (\partial_i h_{ij})^2 u_{(\xi,\epsilon)}^2 \right. \\ & \quad \left. - \int_{\mathbb{R}_+^n} \mu\lambda^{2d} f(\lambda^{-2}|\bar{x}|^2) H_{ij} \partial_i \partial_j u_{(\xi,\epsilon)} w_{(\xi,\epsilon)} \right| \\ & \leq C\mu^{\frac{2(n-1)}{n-2}} \lambda^{\frac{(4d+4)(n-1)}{n-2}} + C\mu\lambda^{2d+2} \left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}} + C\left(\frac{\lambda}{\rho}\right)^{n-2}. \end{aligned}$$

*Proof.* It follows from the definition of  $v_{(\xi, \epsilon)}$  that

$$\begin{aligned} & \int_{\mathbb{R}_+^n} \left\{ \langle dv_{(\xi, \epsilon)}, d(v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \rangle_g + c_n R_g v_{(\xi, \epsilon)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \right\} \\ & - (n-2) \int_{\partial \mathbb{R}_+^n} |v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} v_{(\xi, \epsilon)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) = 0 \end{aligned}$$

Thus,

$$\begin{aligned} & \int_{\mathbb{R}_+^n} \left\{ |dv_{(\xi, \epsilon)}|_g^2 - \langle dv_{(\xi, \epsilon)}, du_{(\xi, \epsilon)} \rangle_g + c_n R_g (v_{(\xi, \epsilon)}^2 - u_{(\xi, \epsilon)} v_{(\xi, \epsilon)}) \right\} \quad (3.5) \\ & - (n-2) \int_{\partial \mathbb{R}_+^n} \left\{ |v_{(\xi, \epsilon)}|^{\frac{2(n-1)}{n-2}} - |v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} v_{(\xi, \epsilon)} u_{(\xi, \epsilon)} \right\} = 0. \end{aligned}$$

We set

$$\begin{aligned} \varrho &= \int_{\mathbb{R}_+^n} \left\{ \langle du_{(\xi, \epsilon)}, d(v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \rangle_g + c_n R_g u_{(\xi, \epsilon)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \right\} \quad (3.6) \\ & - \int_{\mathbb{R}_+^n} h_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) - (n-2) \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{n}{n-2}} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \end{aligned}$$

Thus,

$$\begin{aligned} \varrho &= \int_{\mathbb{R}_+^n} \left\{ -|du_{(\xi, \epsilon)}|_g^2 + \langle du_{(\xi, \epsilon)}, dv_{(\xi, \epsilon)} \rangle_g + c_n R_g (u_{(\xi, \epsilon)} v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}^2) \right\} \\ & - \int_{\mathbb{R}_+^n} h_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) - (n-2) \int_{\partial \mathbb{R}_+^n} \left\{ u_{(\xi, \epsilon)}^{\frac{n}{n-2}} v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}^{\frac{2(n-1)}{n-2}} \right\} \end{aligned}$$

Hence, summing (3.5) and (3.7),

$$\begin{aligned} \varrho &= \int_{\mathbb{R}_+^n} \left\{ |dv_{(\xi, \epsilon)}|_g^2 - |du_{(\xi, \epsilon)}|_g^2 + c_n R_g (v_{(\xi, \epsilon)}^2 - u_{(\xi, \epsilon)}^2) \right\} \quad (3.7) \\ & - \int_{\mathbb{R}_+^n} h_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \\ & - (n-2) \int_{\partial \mathbb{R}_+^n} \left\{ (|v_{(\xi, \epsilon)}|^{\frac{2(n-1)}{n-2}} - u_{(\xi, \epsilon)}^{\frac{2(n-1)}{n-2}}) + (u_{(\xi, \epsilon)}^{\frac{2}{n-2}} - |v_{(\xi, \epsilon)}|^{\frac{2}{n-2}}) u_{(\xi, \epsilon)} v_{(\xi, \epsilon)} \right\}. \end{aligned}$$

Then

$$\begin{aligned} \varrho &= \int_{\mathbb{R}_+^n} \left\{ |dv_{(\xi, \epsilon)}|_g^2 + c_n R_g v_{(\xi, \epsilon)}^2 \right\} - \int_{\partial \mathbb{R}_+^n} \left\{ \frac{(n-2)^2}{n-1} |v_{(\xi, \epsilon)}|^{\frac{2(n-1)}{n-2}} + \frac{n-2}{n-1} u_{(\xi, \epsilon)}^{\frac{2(n-1)}{n-2}} \right\} \quad (3.8) \\ & - \int_{\partial \mathbb{R}_+^n} \left\{ \frac{n-2}{n-1} |v_{(\xi, \epsilon)}|^{\frac{2(n-1)}{n-2}} + \frac{(n-2)^2}{n-1} u_{(\xi, \epsilon)}^{\frac{2(n-1)}{n-2}} \right\} \\ & - (n-2) \int_{\partial \mathbb{R}_+^n} \left( u_{(\xi, \epsilon)}^{\frac{2}{n-2}} - |v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} \right) u_{(\xi, \epsilon)} v_{(\xi, \epsilon)} + 2(n-2) \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{2(n-1)}{n-2}} \\ & - \int_{\mathbb{R}_+^n} \left\{ |du_{(\xi, \epsilon)}|_g^2 + c_n R_g u_{(\xi, \epsilon)}^2 + h_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \right\}. \end{aligned}$$

We set

$$B = \int_{\mathbb{R}_+^n} \left\{ |du_{(\xi, \epsilon)}|_g^2 - |du_{(\xi, \epsilon)}|^2 + c_n R_g u_{(\xi, \epsilon)}^2 + h_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)}) \right\}$$

and observe that  $\int_{\mathbb{R}_+^n} |du_{(\xi, \epsilon)}|^2 = (n-2) \int_{\partial \mathbb{R}_+^n} u_{(\xi, \epsilon)}^{\frac{2(n-1)}{n-2}}$ . Hence,

$$\begin{aligned} \mathcal{F}_g(\xi, \epsilon) - B &= \frac{n-2}{n-1} \int_{\partial \mathbb{R}_+^n} \left\{ |v_{(\xi, \epsilon)}|^{\frac{2(n-1)}{n-2}} - u_{(\xi, \epsilon)}^{\frac{2(n-1)}{n-2}} \right\} \\ &\quad - (n-2) \int_{\partial \mathbb{R}_+^n} \left( |v_{(\xi, \epsilon)}|^{\frac{2}{n-2}} - u_{(\xi, \epsilon)}^{\frac{2}{n-2}} \right) u_{(\xi, \epsilon)} v_{(\xi, \epsilon)} + \varrho \\ &= O \left\{ \lambda^{\frac{(4d+4)(n-1)}{n-2}} \mu^{\frac{2(n-1)}{n-2}} + \left( \frac{\lambda}{\rho} \right)^{n-1} \right\} + \varrho \end{aligned} \quad (3.9)$$

where in the last inequality we used Lemma 3.5.

On the other hand,

$$\begin{aligned} B &= \frac{1}{2} \int_{B_\rho^+(0)} h_{ij} h_{jl} \partial_i u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)} - \frac{c_n}{4} \int_{B_\rho^+(0)} (\partial_i h_{ij})^2 u_{(\xi, \epsilon)}^2 \\ &\quad + \int_{\mathbb{R}_+^n} \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} w_{(\xi, \epsilon)} + e_1 + e_2 + e_3 + e_4 + e_5 \end{aligned} \quad (3.10)$$

where

$$\begin{aligned} e_1 &= - \int_{\mathbb{R}_+^n} h_{ij} \partial_i u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)} + c_n \int_{\mathbb{R}_+^n} \partial_i \partial_j h_{ij} u_{(\xi, \epsilon)}^2, \\ e_2 &= \int_{\mathbb{R}_+^n} (g^{ij} - \delta_{ij} + h_{ij}) \partial_i u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)} - \int_{B_\rho^+(0)} \frac{1}{2} h_{ij} h_{jl} \partial_i u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)} \\ &= \int_{\mathbb{R}_+^n \setminus B_\rho^+(0)} (g^{ij} - \delta_{ij} + h_{ij}) \partial_i u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)} \\ &\quad + \int_{B_\rho^+(0)} \left\{ g^{ij} - \delta_{ij} + h_{ij} - \frac{1}{2} h_{ij} h_{jl} \right\} \partial_i u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)}, \\ e_3 &= c_n \int_{\mathbb{R}_+^n} (R_g - \partial_i \partial_j h_{ij}) u_{(\xi, \epsilon)}^2 + c_n \int_{B_\rho^+(0)} \frac{1}{4} (\partial_i h_{ij})^2 u_{(\xi, \epsilon)}^2 \\ &= c_n \int_{\mathbb{R}_+^n \setminus B_\rho^+(0)} (R_g - \partial_i \partial_j h_{ij}) u_{(\xi, \epsilon)}^2 + c_n \int_{B_\rho^+(0)} \left\{ R_g + \frac{1}{4} (\partial_i h_{ij})^2 \right\} u_{(\xi, \epsilon)}^2, \\ e_4 &= \int_{\mathbb{R}_+^n} h_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} (v_{(\xi, \epsilon)} - u_{(\xi, \epsilon)} - w_{(\xi, \epsilon)}), \\ e_5 &= - \int_{\mathbb{R}_+^n \setminus B_\rho^+(0)} \mu \lambda^{2d} f(\lambda^{-2} |\bar{x}|^2) H_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} w_{(\xi, \epsilon)} + \int_{\mathbb{R}_+^n \setminus B_\rho^+(0)} h_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} w_{(\xi, \epsilon)}. \end{aligned}$$



For the expression of  $e_3$  we used the fact that  $\partial_j h_{ij}(x) = 0$  for  $|x| \leq \rho$ . We are going to use this same fact in the rest of this proof.

Now we are going to estimate the terms  $e_1, \dots, e_5$ . First observe that for  $|x| \leq \rho$  we have

$$\begin{aligned} \left| g^{ij}(x) - \delta_{ij} + h_{ij}(x) - \frac{1}{2} h_{ij} h_{jl}(x) \right| &\leq C|h(x)|^3 \leq C\mu^3(\lambda + |x|)^{6d+6} \\ &\leq C\mu^3(\lambda + |x|)^{\frac{n-1}{n-2}(4d+4)} \end{aligned} \quad (3.11)$$

and

$$\begin{aligned} \left| R_g(x) + \frac{1}{4}(\partial_l h_{ij})^2(x) \right| &\leq C|h(x)|^2|\partial^2 h(x)| + C|h(x)||\partial h(x)|^2 \\ &\leq C\mu^3(\lambda + |x|)^{6d+4} \leq C\mu^3(\lambda + |x|)^{\frac{n-1}{n-2}(4d+4)-2}. \end{aligned} \quad (3.12)$$

Here, we used Lemma 2.1.

From the identity  $u_{(\xi, \epsilon)} \partial_i \partial_j u_{(\xi, \epsilon)} - \frac{n}{n-2} \partial_i u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)} = -\frac{1}{n-2} |du_{(\xi, \epsilon)}|^2 \delta_{ij}$  and the fact that  $\sum_{j=1}^{n-1} h_{jj} = 0$  we see that

$$\begin{aligned} \frac{n}{n-2} \int_{\mathbb{R}_+^n} h_{ij} \partial_i u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)} &= \int_{\mathbb{R}_+^n} h_{ij} u_{(\xi, \epsilon)} \partial_i \partial_j u_{(\xi, \epsilon)} \\ &= - \int_{\mathbb{R}_+^n} \partial_i h_{ij} u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)} - \int_{\mathbb{R}_+^n} h_{ij} \partial_i u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)}, \end{aligned}$$

where in the last equality we integrated by parts. Thus,

$$e_1 = \frac{n-2}{2(n-1)} \int_{\mathbb{R}_+^n \setminus B_\rho^+(0)} \partial_i h_{ij} u_{(\xi, \epsilon)} \partial_j u_{(\xi, \epsilon)} + c_n \int_{\mathbb{R}_+^n \setminus B_\rho^+(0)} \partial_i \partial_j h_{ij} u_{(\xi, \epsilon)}^2. \quad (3.13)$$

Then we use the identities (3.13), (3.11) and (3.12) to estimate  $e_1, e_2$  and  $e_3$

respectively and conclude that

$$\begin{aligned}
|e_1| &\leq C\rho\left(\frac{\lambda}{\rho}\right)^{n-2}, \\
|e_2| &\leq C\left(\frac{\lambda}{\rho}\right)^{n-2} + C\mu^3\lambda^{\frac{n-1}{n-2}(4d+4)}, \\
|e_3| &\leq C\rho^2\left(\frac{\lambda}{\rho}\right)^{n-2} + C\mu^3\lambda^{\frac{n-1}{n-2}(4d+4)} \\
|e_4| &\leq C\int_{\mathbb{R}_+^n} |h|\partial^2 u_{(\xi,\epsilon)}\|v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)} - w_{(\xi,\epsilon)}| \\
&\leq C\|h\partial^2 u_{(\xi,\epsilon)}\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)}\|v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)} - w_{(\xi,\epsilon)}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)} \\
&\leq C\left\{\mu\lambda^{2d+2} + \left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}}\right\}\cdot\left\{\mu^{\frac{n}{n-2}}\lambda^{\frac{(2d+2)n}{n-2}} + \left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}}\right\} \\
&\leq C\mu^{\frac{2(n-1)}{n-2}}\lambda^{\frac{(4d+4)(n-1)}{n-2}} + \mu\lambda^{2d+2}\left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}} + \mu^{\frac{n}{n-2}}\lambda^{\frac{(2d+2)n}{n-2}}\left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}} + \left(\frac{\lambda}{\rho}\right)^{n-2}, \\
|e_5| &\leq \rho^{2d+2}\left(\frac{\lambda}{\rho}\right)^{n-2}.
\end{aligned} \tag{3.14}$$

Now we are going to estimate  $\varrho$  using its definition (equation (3.6)). Integrating by parts and using the second equation of (2.1), we obtain

$$\begin{aligned}
|\varrho| &\leq \int_{\mathbb{R}_+^n} \left| -\Delta_g u_{(\xi,\epsilon)}(v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)}) + c_n R_g u_{(\xi,\epsilon)}(v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)}) \right. \\
&\quad \left. - h_{ij}\partial_i\partial_j u_{(\xi,\epsilon)}(v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)}) \right| \\
&\leq \left\| \Delta_g u_{(\xi,\epsilon)} - c_n R_g u_{(\xi,\epsilon)} + h_{ij}\partial_i\partial_j u_{(\xi,\epsilon)} \right\|_{L^{\frac{2n}{n+2}}(\mathbb{R}_+^n)} \|v_{(\xi,\epsilon)} - u_{(\xi,\epsilon)}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}_+^n)} \\
&\leq C\left\{\mu^2\lambda^{4d+4} + \left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}}\right\}\cdot\left\{\mu\lambda^{2d+2} + \left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}}\right\} \\
&\leq C\mu^3\lambda^{6d+6} + C\mu\lambda^{2d+2}\left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}} + C\left(\frac{\lambda}{\rho}\right)^{n-2}.
\end{aligned} \tag{3.15}$$

Here, we used Proposition 3.1 and Corollary 3.2 in the second inequality.

The result now follows from (3.9), (3.10), (3.14) and (3.15).  $\square$

## 4 Finding a critical point of an auxiliary function

Let us follow the notations of the last section. We define

$$F(\xi, \epsilon) = \frac{1}{2} \int_{\mathbb{R}_+^n} \bar{H}_{il}\bar{H}_{jl}\partial_i u_{(\xi,\epsilon)}\partial_j u_{(\xi,\epsilon)} - \frac{c_n}{4} \int_{\mathbb{R}_+^n} (\partial_l \bar{H}_{ij})^2 u_{(\xi,\epsilon)}^2 + \int_{\mathbb{R}_+^n} \bar{H}_{ij}\partial_i\partial_j u_{(\xi,\epsilon)} z_{(\xi,\epsilon)}$$

where  $z_{(\xi, \epsilon)}$  is the unique element of  $\Sigma_{(\xi, \epsilon)}$  that satisfies

$$\int_{\mathbb{R}_+^n} \langle dz_{(\xi, \epsilon)}, d\psi \rangle - \int_{\partial\mathbb{R}_+^n} nu_{(\xi, \epsilon)}^{\frac{2}{n-2}} z_{(\xi, \epsilon)} \psi = - \int_{\mathbb{R}_+^n} \bar{H}_{ij} \partial_i \partial_j u_{(\xi, \epsilon)} \psi \quad (4.1)$$

for any  $\psi \in \Sigma_{(\xi, \epsilon)}$ . The function  $z_{(\xi, \epsilon)}$  is obtained in Proposition 2.7 with  $h_{ab} = 0$ .

In this section we will show that the function  $F(\xi, \epsilon)$  has a critical point, which is a strict local minimum. Recall that throughout this article we use indices  $1 \leq i, j, k, l, m, p, q, r, s \leq n-1$ .

Since  $\bar{H}_{ab}(-x) = \bar{H}_{ab}(x)$  for any  $x \in \mathbb{R}_+^n$ , the function  $F(\xi, \epsilon)$  satisfies  $F(\xi, \epsilon) = F(-\xi, \epsilon)$  for all  $(\xi, \epsilon) \in \mathbb{R}^{n-1} \times (0, \infty)$ . In particular,

$$\frac{\partial}{\partial \xi_p} F(0, \epsilon) = \frac{\partial^2}{\partial \epsilon \partial \xi_p} F(0, \epsilon) = 0, \quad \text{for all } \epsilon > 0. \quad (4.2)$$

**Proposition 4.1.** *We have*

$$\begin{aligned} \int_{S_r^{n-2}} (\partial_l H_{ij})^2(x) x^p x^q &= \frac{2\sigma_{n-2} r^{n+2}}{(n-1)(n+1)} (W_{ipjl} + W_{iljp}) (W_{iqjl} + W_{iljq}) \\ &\quad + \frac{\sigma_{n-2} r^{n+2}}{(n-1)(n+1)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \end{aligned}$$

and

$$\begin{aligned} \int_{S_r^{n-2}} (H_{ij})^2(x) x^p x^q &= \frac{2\sigma_{n-2} r^{n+4}}{(n-1)(n+1)(n+3)} (W_{ipjl} + W_{iljp}) (W_{iqjl} + W_{iljq}) \\ &\quad + \frac{\sigma_{n-2} r^{n+4}}{2(n-1)(n+1)(n+3)} (W_{ikjl} + W_{iljk})^2 \delta_{pq}. \end{aligned}$$

*Proof.* Observe that

$$\int_{S_r^{n-2}} (\partial_l H_{ij})^2(x) x^p x^q = \int_{S_r^{n-2}} (W_{iljr} + W_{irjl}) (W_{iljm} + W_{imjl}) x^r x^m x^p x^q$$

and

$$\int_{S_r^{n-2}} (H_{ij})^2(x) x^p x^q = \int_{S_r^{n-2}} W_{ikjl} W_{irjm} x^k x^l x^r x^m x^p x^q.$$

Now we just need to apply Corollary A-3 in the Appendix.  $\square$

**Proposition 4.2.** *We have*

$$\begin{aligned} \int_{S_r^{n-2}} (\partial_l \bar{H}_{ij})^2(x) x^p x^q &= \frac{2\sigma_{n-2} r^{n+2}}{(n-1)(n+1)(n+3)} (W_{ipjl} + W_{iljp}) (W_{iqjl} + W_{iljq}) \\ &\quad \cdot \left\{ (n+3) f(r^2)^2 + 8r^2 f(r^2) f'(r^2) + 4r^4 f'(r^2)^2 \right\} \\ &\quad + \frac{\sigma_{n-2} r^{n+2}}{(n-1)(n+1)(n+3)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \\ &\quad \cdot \left\{ (n+3) f(r^2)^2 + 4r^2 f(r^2) f'(r^2) + 2r^4 f'(r^2)^2 \right\}. \end{aligned}$$

*Proof.* Since

$$\partial_l \bar{H}_{ij}(x) = f(|\bar{x}|^2) \partial_l H_{ij}(x) + 2f'(|\bar{x}|^2) x^l H_{ij}(x)$$

we obtain

$$\begin{aligned} (\partial_l \bar{H}_{ij})^2(x) &= f(|\bar{x}|^2)^2 (\partial_l H_{ij})^2(x) + 4f(|\bar{x}|^2) f'(|\bar{x}|^2) x^l \partial_l H_{ij} H_{ij}(x) + 4|\bar{x}|^2 f'(|\bar{x}|^2)^2 (H_{ij})^2 \\ &= f(|\bar{x}|^2)^2 (\partial_l H_{ij})^2(x) + 8f(|\bar{x}|^2) f'(|\bar{x}|^2) (H_{ij})^2(x) + 4|\bar{x}|^2 f'(|\bar{x}|^2)^2 (H_{ij})^2(x) \end{aligned}$$

Hence,

$$\begin{aligned} \int_{S_r^{n-2}} (\partial_l \bar{H}_{ij})^2(x) x^p x^q &= f(|\bar{x}|^2)^2 \int_{S_r^{n-2}} (\partial_l H_{ij})^2(x) x^p x^q \\ &\quad + \left( 8f(|\bar{x}|^2) f'(|\bar{x}|^2) + 4r^2 f'(|\bar{x}|^2)^2 \right) \int_{S_r^{n-2}} (H_{ij})^2(x) x^p x^q \end{aligned}$$

and the result follows from Proposition 4.1.  $\square$

**Corollary 4.3.** *We have*

$$\begin{aligned} \int_{S_r^{n-2}} (\partial_l \bar{H}_{ij})^2(x) &= \\ &= \frac{\sigma_{n-2} r^n}{(n-1)(n+1)} (W_{ikjl} + W_{iljk})^2 \left\{ (n+1) f(r^2)^2 + 4r^2 f(r^2) f'(r^2) + 2r^4 f'(r^2)^2 \right\}. \end{aligned}$$

*Proof.* By Proposition 4.2,

$$\begin{aligned} r^2 \int_{S_r^{n-2}} (\partial_l \bar{H}_{ij})^2(x) &= \sum_{p=1}^{n-1} \int_{S_r^{n-2}} (\partial_l \bar{H}_{ij})^2(x) (x^p)^2 \\ &= \frac{2\sigma_{n-2} r^{n+2}}{(n-1)(n+1)(n+3)} (W_{ikjl} + W_{iljk})^2 \left\{ (n+3) f(r^2)^2 + 8r^2 f(r^2) f'(r^2) + 4r^4 f'(r^2)^2 \right\} \\ &\quad + \frac{\sigma_{n-2} r^{n+2}}{(n-1)(n+1)(n+3)} (n-1) (W_{ikjl} + W_{iljk})^2 \\ &\quad \cdot \left\{ (n+3) f(r^2)^2 + 4r^2 f(r^2) f'(r^2) + 2r^4 f'(r^2)^2 \right\} \\ &= \frac{\sigma_{n-2} r^{n+2}}{(n-1)(n+1)} (W_{ikjl} + W_{iljk})^2 \left\{ (n+1) f(r^2)^2 + 4r^2 f(r^2) f'(r^2) + 2r^4 f'(r^2)^2 \right\}. \end{aligned}$$

$\square$

**Proposition 4.4.** *We have*

$$\begin{aligned} F(0, \epsilon) &= -\frac{c_n \cdot \sigma_{n-2}}{4(n-1)(n+1)} (W_{ikjl} + W_{iljk})^2 \\ &\cdot \int_0^\infty \int_0^\infty r^n \left\{ (n+1) f(r^2)^2 + 4r^2 f(r^2) f'(r^2) + 2r^4 f'(r^2)^2 \right\} e^{n-2} ((\epsilon+t)^2 + r^2)^{2-n} dr dt. \end{aligned}$$

*Proof.* It follows from symmetry arguments that  $z_{(0,\epsilon)} = 0$  and

$$\begin{aligned} & \int_{S^{n-2}} \bar{H}_{ij} \bar{H}_{jl} \partial_i u_{(0,\epsilon)} \partial_j u_{(0,\epsilon)}(x) \\ &= \int_{S^{n-2}} \frac{(n-2)^2 \epsilon^{n-2}}{((\epsilon + x_n)^2 + |\bar{x}|^2)^n} f(|\bar{x}|^2)^2 W_{iplq} W_{jrlm} x^i x^j x^p x^q x^r x^m = 0. \end{aligned}$$

Hence, we have

$$\begin{aligned} F(0, \epsilon) &= -\frac{c_n}{4} \int_{\mathbb{R}_+^n} (\partial_l \bar{H}_{ij})^2(x) u_{(0,\epsilon)}^2(x) \\ &= -\frac{c_n}{4} \int_0^\infty \int_0^\infty \int_{S^{n-2}} (\partial_l \bar{H}_{ij})^2(x) u_{(0,\epsilon)}^2(x) d\sigma_r(x) dr dx_n. \end{aligned}$$

The result now follows from Corollary 4.3. □

We write

$$F(0, \epsilon) = -\beta_n \cdot \sum_{q=0}^{2d} \alpha_q \int_0^\infty \int_0^\infty r^{2q+n} \epsilon^{n-2} ((\epsilon + t)^2 + r^2)^{2-n} dr dt,$$

where

$$\beta_n = \frac{c_n \cdot \sigma_{n-2}}{4(n-1)(n+1)} (W_{ikjl} + W_{iljk})^2,$$

and define the coefficients  $\alpha_q \in \mathbb{R}$  by the formula

$$\sum_{q=0}^{2d} \alpha_q s^q = (n+1)f(s)^2 + 4sf(s)f'(s) + 2s^2f'(s)^2. \quad (4.3)$$

Here,  $d$  is the integer in the formula (3.1). Changing variables  $t' = t/\epsilon$  and  $r' = r/\epsilon$  we obtain

$$F(0, \epsilon) = -\beta_n \cdot \sum_{q=0}^{2d} \alpha_q \epsilon^{2q+4} \int_0^\infty \int_0^\infty \frac{r'^{2q+n}}{((1+t')^2 + r'^2)^{n-2}} dr' dt'$$

and, changing variables  $r' = r/(1+t)$ ,

$$F(0, \epsilon) = -\beta_n \cdot \sum_{q=0}^{2d} \alpha_q \epsilon^{2q+4} \int_0^\infty \frac{1}{(1+t)^{n-5-2q}} dt \cdot \int_0^\infty \frac{r'^{2q+n}}{(1+r')^{n-2}} dr'$$

Now, we have

$$\int_0^\infty \frac{1}{(1+t)^{n-5-2q}} dt = \frac{1}{n-6-2q}$$

and

$$\int_0^\infty \frac{r^{2q+n}}{(1+r^2)^{n-2}} dr = \left\{ \prod_{j=0}^q \frac{n-1+2j}{n-5-2j} \right\} \cdot \int_0^\infty \frac{r^{n-2}}{(1+r^2)^{n-2}} dr,$$

where we used Lemma A-1. Hence, we can write

$$F(0, \epsilon) = -\beta_n \cdot I(\epsilon^2) \cdot \int_{r=0}^\infty \frac{r^{n-2}}{(1+r^2)^{n-2}} dr \quad (4.4)$$

where

$$I(s) = \sum_{q=0}^{2d} \frac{\alpha_q}{n-6-2q} \left\{ \prod_{j=0}^q \frac{n-1+2j}{n-5-2j} \right\} s^{q+2}. \quad (4.5)$$

We will now turn our attention to the second order derivatives of the function  $F(\xi, \epsilon)$ .

**Proposition 4.5.** *We have*

$$\begin{aligned} \frac{\partial^2}{\partial \xi_p \partial \xi_q} F(0, \epsilon) &= (n-2)^2 \int_{\mathbb{R}_+^n} \frac{\epsilon^{n-2}}{((\epsilon+x_n)^2 + |\bar{x}|^2)^n} \bar{H}_{pl}(x) \bar{H}_{ql}(x) \\ &\quad - \frac{(n-2)^2}{4} \int_{\mathbb{R}_+^n} \frac{\epsilon^{n-2}}{((\epsilon+x_n)^2 + |\bar{x}|^2)^n} (\partial_l \bar{H}_{ij}(x))^2 x^p x^q \\ &\quad + \frac{(n-2)^2}{8(n-1)} \int_{\mathbb{R}_+^n} \frac{\epsilon^{n-2}}{((\epsilon+x_n)^2 + |\bar{x}|^2)^{n-1}} (\partial_l \bar{H}_{ij}(x))^2 \delta_{pq}. \end{aligned}$$

*Proof.* The proof is the same of Proposition 21 of [6].  $\square$

**Proposition 4.6.** *We have*

$$\begin{aligned} &\frac{\partial^2}{\partial \xi_p \partial \xi_q} F(0, \epsilon) \quad (4.6) \\ &= -\frac{2(n-2)^2 \sigma_{n-2}}{(n-1)(n+1)(n+3)} (W_{ipjl} + W_{iljp})(W_{iqjl} + W_{iljq}) \\ &\quad \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon+x_n)^2 + r^2)^n} r^{n+4} (2f(r^2)f'(r^2) + r^2 f'(r^2)^2) dr dt \\ &\quad - \frac{(n-2)^2 \sigma_{n-2}}{2(n-1)(n+1)(n+3)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \\ &\quad \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon+x_n)^2 + r^2)^n} r^{n+4} \{2f(r^2)f'(r^2) + r^2 f'(r^2)^2\} dr dt \\ &\quad + \frac{(n-2)^2 \sigma_{n-2}}{4(n-1)^2(n+1)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \\ &\quad \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon+x_n)^2 + r^2)^{n-1}} r^{n+4} f'(r^2)^2 dr dt. \end{aligned}$$

*Proof.* It follows from Corollary A-3 in the Appendix that

$$\begin{aligned} \int_{S^{n-2}} \bar{H}_{pl} \bar{H}_{ql}(x) &= \int_{S^{n-2}} f(r^2)^2 H_{pl} H_{ql}(x) = f(r^2)^2 \int_{S^{n-2}} W_{ipkl} W_{jqml} x^i x^j x^k x^m \\ &= \frac{\sigma_{n-2}}{2(n-1)(n+1)} (W_{ipkl} + W_{ilkp})(W_{iqkl} + W_{ilkq}) r^{n+2} f(r^2)^2. \end{aligned}$$

Hence,

$$\begin{aligned} \int_{\mathbb{R}_+^n} \frac{\epsilon^{n-2} \bar{H}_{pl} \bar{H}_{ql}(x)}{((\epsilon + x_n)^2 + |\bar{x}|^2)^n} &= \frac{\sigma_{n-2}}{2(n-1)(n+1)} (W_{ipkl} + W_{ilkp})(W_{iqkl} + W_{ilkq}) \\ &\cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2} r^{n+2}}{((\epsilon + x_n)^2 + r^2)^n} f(r^2)^2 dt dr. \end{aligned} \quad (4.7)$$

It follows from Proposition 4.2 that

$$\begin{aligned} \int_{\mathbb{R}_+^n} \frac{\epsilon^{n-2} (\partial_l \bar{H}_{ij})^2(x) x^p x^q}{((\epsilon + x_n)^2 + |\bar{x}|^2)^n} & \quad (4.8) \\ &= \frac{2\sigma_{n-2}}{(n-1)(n+1)(n+3)} (W_{ipjl} + W_{iljp})(W_{iqjl} + W_{iljq}) \\ &\cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2} r^{n+2}}{((\epsilon + x_n)^2 + r^2)^n} \left\{ (n+3)f(r^2)^2 + 8r^2 f(r^2) f'(r^2) + 4r^4 f'(r^2)^2 \right\} dt dr \\ &+ \frac{\sigma_{n-2}}{(n-1)(n+1)(n+3)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \\ &\cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2} r^{n+2}}{((\epsilon + x_n)^2 + r^2)^n} \left\{ (n+3)f(r^2)^2 + 4r^2 f(r^2) f'(r^2) + 2r^4 f'(r^2)^2 \right\} dt dr. \end{aligned}$$

and from Corollary 4.3 that

$$\begin{aligned} \int_{\mathbb{R}_+^n} \frac{\epsilon^{n-2} (\partial_l \bar{H}_{ij})^2(x) \delta_{pq}}{((\epsilon + x_n)^2 + |\bar{x}|^2)^{n-1}} &= \frac{\sigma_{n-2}}{(n-1)(n+1)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \\ &\cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2} r^n}{((\epsilon + x_n)^2 + r^2)^{n-1}} \left\{ (n+1)f(r^2)^2 + 4r^2 f(r^2) f'(r^2) + 2r^4 f'(r^2)^2 \right\} dt dr. \end{aligned} \quad (4.9)$$

Observe that

$$\begin{aligned} \frac{r^n}{((\epsilon + x_n)^2 + r^2)^{n-1}} \left\{ (n+1)f(r^2)^2 + 4r^2 f(r^2) f'(r^2) \right\} & \quad (4.10) \\ &= \frac{2(n-1)r^{n+2} f(r^2)^2}{((\epsilon + x_n)^2 + r^2)^n} + \frac{d}{dr} \left\{ \frac{r^{n+1} f(r^2)^2}{((\epsilon + x_n)^2 + r^2)^{n-1}} \right\}. \end{aligned}$$

Substituting the equation (4.10) in the equation (4.9) we obtain

$$\begin{aligned}
& \int_{\mathbb{R}_+^n} \frac{\epsilon^{n-2} (\partial_l \bar{H}_{ij})^2(x) \delta_{pq}}{((\epsilon + x_n)^2 + |\bar{x}|^2)^{n-1}} \quad (4.11) \\
&= \frac{2\sigma_{n-2}}{n+1} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2} r^{n+2} f(r^2)^2}{((\epsilon + x_n)^2 + r^2)^n} dt dr \\
&+ \frac{2\sigma_{n-2}}{(n-1)(n+1)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2} r^{n+4} f'(r^2)^2}{((\epsilon + x_n)^2 + r^2)^{n-1}} dt dr,
\end{aligned}$$

since we are assuming that  $n > 4d + 6$ . Now, using the equations (4.7), (4.8) and (4.11) in Proposition 4.5, we obtain

$$\begin{aligned}
& \frac{\partial^2}{\partial \xi_p \partial \xi_q} F(0, \epsilon) \\
&= \frac{(n-2)^2 \sigma_{n-2}}{2(n-1)(n+1)} (W_{ipjl} + W_{iljp})(W_{iqjl} + W_{iljq}) \\
&\quad \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon + x_n)^2 + r^2)^n} r^{n+2} f(r^2)^2 dr dt \\
&- \frac{(n-2)^2 \sigma_{n-2}}{2(n-1)(n+1)(n+3)} (W_{ipjl} + W_{iljp})(W_{iqjl} + W_{iljq}) \\
&\quad \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon + x_n)^2 + r^2)^n} r^{n+2} \left\{ (n+3) f(r^2)^2 + 8r^2 f(r^2) f'(r^2) + 4r^4 f'(r^2)^2 \right\} dr dt \\
&- \frac{(n-2)^2 \sigma_{n-2}}{4(n-1)(n+1)(n+3)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \\
&\quad \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon + x_n)^2 + r^2)^n} r^{n+2} \left\{ (n+3) f(r^2)^2 + 4r^2 f(r^2) f'(r^2) + 2r^4 f'(r^2)^2 \right\} dr dt \\
&+ \frac{(n-2)^2 \sigma_{n-2}}{4(n-1)(n+1)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \\
&\quad \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon + x_n)^2 + r^2)^n} r^{n+2} f(r^2)^2 dr dt \\
&+ \frac{(n-2)^2 \sigma_{n-2}}{4(n-1)^2(n+1)} (W_{ikjl} + W_{iljk})^2 \delta_{pq} \\
&\quad \cdot \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon + x_n)^2 + r^2)^{n-1}} r^{n+4} f'(r^2)^2 dr dt
\end{aligned}$$

and the result follows after we cancel out some terms in the above equation.  $\square$

Let us define constants  $\beta_q$ , for  $q = 0, \dots, 2d-1$ , by the following expression:

$$\sum_{q=0}^{2d-1} \beta_q s^q = 2f(s)f'(s) + s f'(s)^2.$$



**Proposition 4.7.** *We have*

$$\begin{aligned} & \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon + x_n)^2 + r^2)^n} r^{n+4} (2f(r^2)f'(r^2) + r^2 f'(r^2)^2) dr dt \\ &= J(\epsilon^2) \cdot \int_0^\infty \frac{r^{n+2}}{(1+r^2)^n} dr, \end{aligned} \quad (4.12)$$

where

$$J(s) = \sum_{q=0}^{2d-1} \frac{\beta_q s^{q+2}}{n-6-2q} \cdot \left\{ \prod_{j=0}^q \frac{n+3+2j}{n-5-2j} \right\}.$$

*Proof.*

$$\begin{aligned} & \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2}}{((\epsilon + x_n)^2 + r^2)^n} r^{n+4} (2f(r^2)f'(r^2) + r^2 f'(r^2)^2) dr dt \\ &= \sum_{q=0}^{2d-1} \beta_q \int_0^\infty \int_0^\infty \frac{\epsilon^{n-2} r^{n+4+2q}}{((\epsilon + x_n)^2 + r^2)^n} dr dt \\ &= \sum_{q=0}^{2d-1} \beta_q \epsilon^{2q+4} \int_0^\infty \int_0^\infty \frac{r^{n+4+2q}}{((1+x_n)^2 + r^2)^n} dr dt \\ &= \sum_{q=0}^{2d-1} \beta_q \epsilon^{2q+4} \int_0^\infty \frac{1}{(1+t)^{n-5-2q}} dt \int_0^\infty \frac{r^{n+4+2q}}{(1+r^2)^n} dr \end{aligned}$$

Now we observe that

$$\int_0^\infty \frac{1}{(1+t)^{n-5-2q}} dt = \frac{1}{n-6-2q}$$

and apply Lemma A-1 to see that

$$\int_0^\infty \frac{r^{n+4+2q}}{(1+r^2)^n} dr = \left\{ \prod_{j=0}^q \frac{n+3+2j}{n-5-2j} \right\} \cdot \int_0^\infty \frac{r^{n+2}}{(1+r^2)^n} dr.$$

□

#### 4.1 The case $n \geq 53$

In this case we choose  $d = 1$  in the equation (3.1). Then the coefficients  $\alpha_q$  in the equation (4.3) are given by

$$\alpha_0 = (n+1)a_0^2, \quad \alpha_1 = 2(n+3)a_0 a_1, \quad \alpha_2 = (n+7)a_1^2.$$

Thus, derivating  $I(s)$  in the expression (4.5) we obtain

$$\begin{aligned} I'(s) &= \sum_{q=0}^2 \frac{(q+2)\alpha_q}{n-6-2q} \left\{ \prod_{j=0}^q \frac{n-1+2j}{n-5-2j} \right\} s^{q+1} \\ &= \frac{2\alpha_0(n-1)}{(n-6)(n-5)} \cdot s + \frac{3\alpha_1(n-1)(n+1)}{(n-8)(n-5)(n-7)} \cdot s^2 + \frac{4\alpha_2(n-1)(n+1)(n+3)}{(n-10)(n-5)(n-7)(n-9)} \cdot s^3 \\ &= \frac{2(n+1)(n-1)}{n-5} \left\{ \frac{1}{n-6} a_0^2 s + \frac{3(n+3)}{(n-8)(n-7)} a_0 a_1 s^2 + \frac{2(n+3)(n+7)}{(n-10)(n-7)(n-9)} a_1^2 s^3 \right\}. \end{aligned}$$

Now we choose  $a_1 = -1$  and define the polynomial  $p_n$  by

$$p_n(a_0) = \frac{a_0^2}{n-6} - \frac{3(n+3)a_0}{(n-8)(n-7)} + \frac{2(n+3)(n+7)}{(n-10)(n-7)(n-9)}.$$

Hence,

$$I'(1) = \frac{2(n+1)(n-1)}{n-5} p_n(a_0).$$

The discriminant of  $p_n$  is then given by

$$\begin{aligned} \text{discrim}(p_n) &= \frac{(n+3)^2}{(n-7)^2(n-8)^2} \left\{ 9 - \frac{8(n-7)(n-8)^2(n+7)}{(n+3)(n-6)(n-9)(n-10)} \right\} \\ &= \frac{(n+3)^2}{(n-7)^2(n-8)^2} \frac{q(n)}{(n+3)(n-6)(n-9)(n-10)}, \end{aligned}$$

where

$$q(n) = 9(n+3)(n-6)(n-9)(n-10) - 8(n-7)(n-8)^2(n+7).$$

Observe that

$$q'(n) = 4n^3 - 210n^2 + 2082n - 5624$$

and

$$q''(n) = 6(2n^2 - 70n + 347).$$

Since the roots  $\frac{70 \pm \sqrt{2124}}{4}$  of  $q''$  are less than 53, we see that  $q''(n) > 0$  for  $n \geq 53$ . Since  $q(53) = 105696$  and  $q'(53) = 110340$ , we conclude that  $\text{discrim}(p_n) > 0$  for  $n \geq 53$ . Hence, if we set

$$a_0 = \frac{(n+3)(n-6)}{2(n-7)(n-8)} \left\{ 3 + \sqrt{9 - \frac{8(n-7)(n-8)^2(n+7)}{(n+3)(n-6)(n-9)(n-10)}} \right\},$$

then  $s = 1$  is critical point of  $I(s)$ . According to Proposition B-1 in the Appendix,  $I''(1) < 0$  for  $n \geq 53$ .

Now we will handle  $J(s)$ , as defined in Proposition 4.7. We have

$$J(s) = \frac{(n+3)\beta_0 s^2}{(n-6)(n-5)} + \frac{(n+3)(n+5)\beta_1 s^3}{(n-8)(n-5)(n-7)}$$

where

$$\beta_0 = 2a_0a_1 \quad \text{and} \quad \beta_1 = 3a_1^2.$$

Hence,

$$J(s) = \frac{(n+3)a_1}{n-5} \left\{ \frac{2a_0s^2}{n-6} + \frac{3(n+5)a_1s^3}{(n-8)(n-7)} \right\}.$$

If we set  $a_0$  and  $a_1$  as above we have

$$J(1) = \frac{n+3}{(n-8)(n-5)(n-7)} \cdot \left\{ 6 - (n+3) \sqrt{9 - \frac{8(n-7)(n-8)^2(n+7)}{(n+3)(n-6)(n-9)(n-10)}} \right\}$$

According to Proposition B-2 in the Appendix,  $J(1) < 0$  for  $n \geq 53$ .

From the equations (4.2), (4.4), (4.6) and (4.12) and the above results we can conclude the following:

**Proposition 4.8.** *Suppose that  $n \geq 53$ . If we set  $a_1 = -1$  and*

$$a_0 = \frac{(n+3)(n-6)}{2(n-7)(n-8)} \left\{ 3 + \sqrt{9 - \frac{8(n-7)(n-8)^2(n+7)}{(n+3)(n-6)(n-9)(n-10)}} \right\},$$

*then  $I'(1) = 0$ ,  $I''(1) < 0$  and  $J(1) < 0$ . In particular, the function  $F(\xi, \epsilon)$  has a strict local minimum at the point  $(0, 1)$ .*

## 4.2 The case $25 \leq n \leq 52$

In this case we choose  $d = 4$  in the equation (3.1). The coefficients  $\alpha_q$  in the equation (4.3) are then given by

$$\begin{aligned} \alpha_0 &= (n+1)a_0^2, \\ \alpha_1 &= 2(n+3)a_0a_1, \\ \alpha_2 &= 2(n+5)a_0a_2 + (n+7)a_1^2, \\ \alpha_3 &= 2(n+11)a_1a_2 + 2(n+7)a_0a_3, \\ \alpha_4 &= 2(n+15)a_1a_3 + (n+17)a_2^2 + 2(n+9)a_0a_4, \\ \alpha_5 &= 2(n+23)a_2a_3 + 2(n+19)a_1a_4, \\ \alpha_6 &= (n+31)a_3^2 + 2(n+29)a_2a_4, \\ \alpha_7 &= 2(n+39)a_3a_4, \\ \alpha_8 &= (n+49)a_4^2. \end{aligned}$$

Thus, derivating  $I(s)$  in the expression (4.5) we obtain

$$I'(s) = \sum_{q=0}^8 \frac{(q+2)\alpha_q}{n-6-2q} \left\{ \prod_{j=0}^q \frac{n-1+2j}{n-5-2j} \right\} s^{q+1}.$$

Now we choose  $a_1 = -3/5$ ,  $a_2 = 1/8$ ,  $a_3 = -1/125$ ,  $a_4 = 10^{-4}$  and define the polynomial  $r_n$  by  $r_n(a_0) = I'(1)$ . Hence,

$$r_n(a_0) = \frac{2(n-1)(n+1)}{(n-6)(n-5)} \cdot a_0^2 + \left\{ \sum_{q=1}^4 \gamma_q(n) \frac{q+2}{n-6-2q} \prod_{j=0}^q \frac{n-1+2j}{n-5-2j} \right\} \cdot a_0 \\ + \sum_{q=2}^8 \delta_q(n) \frac{q+2}{n-6-2q} \prod_{j=0}^q \frac{n-1+2j}{n-5-2j},$$

where

$$\gamma_1(n) = -\frac{6}{5}(n+3), \quad \gamma_2(n) = \frac{n+5}{4}, \quad \gamma_3(n) = -\frac{2}{125}(n+7), \quad \gamma_4(n) = \frac{n+9}{5000}, \\ \delta_2(n) = \frac{9(n+7)}{25}, \delta_3(n) = -\frac{3(n+11)}{20}, \delta_4(n) = \frac{1009n+16385}{40000}, \delta_5(n) = -\frac{53n+1207}{25000}, \\ \delta_6(n) = \frac{89n+2709}{10^6}, \quad \delta_7(n) = -\frac{n+39}{625000}, \quad \delta_8(n) = \frac{n+49}{10^8}.$$

Direct computations show that  $\text{discrim}(r_n) > 0$  for  $25 \leq n \leq 52$ .

If we choose

$$a_0 = \frac{(n-6)(n-5)}{4(n-1)(n+1)} \cdot \left\{ -\sum_{q=1}^4 \gamma_q(n) \frac{q+2}{n-6-2q} \prod_{j=0}^q \frac{n-1+2j}{n-5-2j} + \sqrt{\text{discrim}(r_n)} \right\}$$

then  $s = 1$  is critical point of  $I(s)$ . For  $25 \leq n \leq 52$ , direct computations show that  $I''(1)$  is of the form  $-e_1 - e_2 \sqrt{e_3}$ , where  $e_1, e_2, e_3$  are positive rational numbers.

The function  $J(s)$ , defined in Proposition 4.7, is written as

$$J(s) = \sum_{q=0}^7 \frac{\beta_q s^{q+2}}{n-6-2q} \cdot \left\{ \prod_{j=0}^q \frac{n+3+2j}{n-5-2j} \right\}.$$

where

$$\beta_0 = 2a_0 a_1, \quad \beta_1 = 4a_0 a_2 + 3a_1^2, \quad \beta_2 = 6a_0 a_3 + 10a_1 a_2, \quad \beta_3 = 8a_0 a_4 + 14a_1 a_3 + 8a_2^2, \\ \beta_4 = 18a_1 a_4 + 22a_2 a_3, \quad \beta_5 = 28a_2 a_4 + 15a_3^2, \quad \beta_6 = 38a_3 a_4, \quad \beta_7 = 24a_4^2.$$

For  $25 \leq n \leq 52$ , direct computations show that  $J(1)$  is of the form  $-e_1 - e_2 \sqrt{e_3}$ , where  $e_1, e_2, e_3$  are positive rational numbers. From the equations (4.2), (4.4), (4.6) and (4.12) and the above results we can conclude the following:

**Proposition 4.9.** *Suppose that  $25 \leq n \leq 52$ . If  $a_1 = -3/5$ ,  $a_2 = 1/8$ ,  $a_3 = -1/125$ ,  $a_4 = 10^{-4}$  and*

$$a_0 = \frac{(n-6)(n-5)}{4(n-1)(n+1)} \cdot \left\{ -\sum_{q=1}^4 \gamma_q(n) \frac{q+2}{n-6-2q} \prod_{j=0}^q \frac{n-1+2j}{n-5-2j} + \sqrt{\text{discrim}(r_n)} \right\}$$

then  $I'(1) = 0$ ,  $I''(1) < 0$  and  $J(1) < 0$ . In particular, the function  $F(\xi, \epsilon)$  has a strict local minimum at the point  $(0, 1)$ .

## 5 Proof of the main theorem

In this section we will make use of the two-tensor  $H$ , defined on  $\mathbb{R}_+^n$ , the polynomial  $f$  and the open set  $\Omega \subset \mathbb{R}^{n-1} \times (0, \infty)$ , which were defined in Section 3. As in Sections 4.1 and 4.2, we fix  $d = 1$  if  $n \geq 53$  and  $d = 4$  if  $25 \leq n \leq 52$ . We set  $D_r(0) = \{x \in \partial\mathbb{R}_+^n; |x| < r\}$ .

The basic ingredient in the proof of the Main Theorem is the following result:

**Proposition 5.1.** *Assume that  $n \geq 25$ . Let  $g$  be a smooth Riemannian metric on  $\mathbb{R}_+^n$  expressed as  $g = \exp(h)$ , where  $h$  is a symmetric trace-free two-tensor on  $\mathbb{R}_+^n$  satisfying the following properties:*

$$\begin{cases} h_{ab}(x) = \mu \lambda^{2d} f(\lambda^{-2}|\bar{x}|) H_{ab}(x), & \text{for } |x| \leq \rho, \\ h_{ab}(x) = 0, & \text{for } |x| \geq 1, \\ h_{nb}(x) = 0, & \text{for } x \in \mathbb{R}_+^n, \\ \partial_n h_{ab}(x) = 0, & \text{for } x \in \partial\mathbb{R}_+^n, \end{cases} \quad (5.1)$$

where  $a, b = 1, \dots, n$ . We also assume that

$$|h(x)| + |\partial h(x)| + |\partial^2 h(x)| \leq \alpha \leq \alpha_1, \quad \text{for all } x \in \mathbb{R}_+^n,$$

where  $\alpha_1$  is the constant obtained in Proposition 2.8.

If  $\alpha$  and  $\mu^{-2} \lambda^{n-4d-6} \rho^{2-n}$  are sufficiently small, then there exists a positive smooth function  $v$  satisfying

$$\begin{cases} \Delta_g v - c_n R_g v = 0, & \text{in } \mathbb{R}_+^n, \\ \frac{\partial}{\partial x_n} v - d_n \kappa_g v + (n-2)v^{\frac{n}{n-2}} = 0, & \text{on } \partial\mathbb{R}_+^n \end{cases} \quad (5.2)$$

and

$$\int_{\partial\mathbb{R}_+^n} v^{\frac{2(n-1)}{n-2}} < \left( \frac{Q(B^n, \partial B)}{n-2} \right)^{n-1}. \quad (5.3)$$

Moreover, there exists  $c = c(n) > 0$  such that

$$\sup_{D_\lambda(0)} v \geq c \lambda^{\frac{2-n}{2}}. \quad (5.4)$$

*Proof.* It follows from the fact that

$$(n+1)f(s)^2 + 4sf(s)f'(s) + 2s^2f'(s)^2 = (n-1)f(s)^2 + 2(f(s) + sf'(s))^2$$

and Proposition 4.4 that  $F(0, 1) < 0$ . According to Propositions 4.8 and 4.9, we can choose the coefficients  $a_0, \dots, a_d$  in the formula (3.1) such that the point  $(0, 1)$  is a strict local minimum of  $F$ . Hence, we can find an open set  $\Omega' \subset \Omega$  such that  $(0, 1) \in \Omega'$  and

$$F(0, 1) < \inf_{(\xi, \epsilon) \in \partial\Omega'} F(\xi, \epsilon) < 0.$$

Observe that  $u_{(\lambda\xi, \lambda\epsilon)}(\lambda x) = \lambda^{-\frac{n-2}{2}} u_{(\xi, \epsilon)}(x)$  and  $w_{(\lambda\xi, \lambda\epsilon)}(\lambda x) = \mu \lambda^{2d+2-\frac{n-2}{2}} z_{(\xi, \epsilon)}(x)$  for all  $x \in \mathbb{R}_+^n$ . Here,  $w_{(\xi, \epsilon)}$  and  $z_{(\xi, \epsilon)}$  are the functions defined by the formulas (3.2) and (4.1) respectively. Thus, it follows from Proposition 3.6 that

$$|\mathcal{F}_g(\lambda\xi, \lambda\epsilon) - \mu^2 \lambda^{4d+4} F(\xi, \epsilon)| \leq C \mu^{\frac{2(n-1)}{n-2}} \lambda^{\frac{(4d+4)(n-1)}{n-2}} + C \mu \lambda^{2d+2} \left(\frac{\lambda}{\rho}\right)^{\frac{n-2}{2}} + C \left(\frac{\lambda}{\rho}\right)^{n-2}$$

for all  $(\xi, \epsilon) \in \Omega$ . Hence,

$$\begin{aligned} \left| \mu^{-2} \lambda^{-4d-4} \mathcal{F}_g(\lambda\xi, \lambda\epsilon) - F(\xi, \epsilon) \right| &\leq C \mu^{\frac{2}{n-2}} \lambda^{\frac{4d+4}{n-2}} \\ &\quad + C \mu^{-1} \lambda^{\frac{n-4d-6}{2}} \rho^{\frac{2-n}{2}} + C \mu^{-2} \lambda^{n-4d-6} \rho^{2-n} \end{aligned}$$

for all  $(\xi, \epsilon) \in \Omega$ . If  $\mu^{-2} \lambda^{n-4d-6} \rho^{2-n}$  is sufficiently small then we have

$$\mathcal{F}_g(0, \lambda) < \inf_{(\xi, \epsilon) \in \partial\Omega'} \mathcal{F}_g(\lambda\xi, \lambda\epsilon) < 0.$$

Thus we conclude that there exists a point  $(\bar{\xi}, \bar{\epsilon}) \in \Omega'$  such that

$$\mathcal{F}_g(\lambda\bar{\xi}, \lambda\bar{\epsilon}) = \inf_{(\xi, \epsilon) \in \Omega'} \mathcal{F}_g(\lambda\xi, \lambda\epsilon) < 0.$$

By Proposition 2.9, the function  $v = v_{(\lambda\bar{\xi}, \lambda\bar{\epsilon})}$  obtained in Proposition 2.8 is a positive smooth solution to the equations (5.2). Hence, by the definition of  $\mathcal{F}_g$  (see the formula (2.19)) and the formula (2.2), we have

$$\frac{n-2}{n-1} \int_{\partial\mathbb{R}_+^n} v^{\frac{2(n-1)}{n-2}} = \frac{n-2}{n-1} \left( \frac{Q(B^n, \partial B)}{n-2} \right)^{n-1} + \mathcal{F}(\lambda\bar{\xi}, \lambda\bar{\epsilon}).$$

This implies the inequality (5.3).

In order to prove the inequality (5.4), observe that

$$\|v - u_{(\lambda\bar{\xi}, \lambda\bar{\epsilon})}\|_{L^{\frac{2(n-1)}{n-2}}(D_\lambda(0))} \leq \|v - u_{(\lambda\bar{\xi}, \lambda\bar{\epsilon})}\|_{L^{\frac{2(n-1)}{n-2}}(\partial\mathbb{R}_+^n)} \leq C\alpha$$

by Propositions 2.2 and 2.8. Hence,

$$|D_\lambda(0)|^{\frac{n-2}{2(n-1)}} \sup_{D_\lambda(0)} v \geq \|v\|_{L^{\frac{2(n-1)}{n-2}}(D_\lambda(0))} \geq -C\alpha + \|u_{(\lambda\bar{\xi}, \lambda\bar{\epsilon})}\|_{L^{\frac{2(n-1)}{n-2}}(D_\lambda(0))}.$$

Now, the inequality (5.4) follows from choosing  $\alpha$  sufficiently small.  $\square$

Now the Main Theorem follows from the next theorem, using the conformal equivalence between  $B^n \setminus \{(0, \dots, 0, -1)\}$  and  $\mathbb{R}_+^n$  (see Lemma 2.3), the properties (2.8) and Lemma 2.10.

**Theorem 5.2.** *Assume that  $n \geq 25$ . Then there exists a smooth Riemannian metric  $g$  on  $\mathbb{R}_+^n$  with the following properties:*

(a)  $g_{ab}(x) = \delta_{ab}$  for  $|x| \geq 1/2$ ;

- (b)  $g$  is not conformally flat;  
(c)  $\partial\mathbb{R}_+^n$  is totally geodesic with respect to the induced metric by  $g$ ;  
(d) there exists a sequence of positive smooth functions  $\{v_\nu\}_{\nu=1}^\infty$  satisfying

$$\begin{cases} \Delta_g v_\nu - c_n R_g v_\nu = 0, & \text{in } \mathbb{R}_+^n, \\ \frac{\partial}{\partial x_n} v_\nu - d_n \kappa_g v_\nu + (n-2)v_\nu^{\frac{n}{n-2}} = 0, & \text{on } \partial\mathbb{R}_+^n, \end{cases} \quad (5.5)$$

for all  $\nu$ ,

$$\int_{\partial\mathbb{R}_+^n} v_\nu^{\frac{2(n-1)}{n-2}} < \left( \frac{Q(B^n, \partial B)}{n-2} \right)^{n-1},$$

for all  $\nu$ , and  $\sup_{D_1(0)} v_\nu \rightarrow \infty$  as  $\nu \rightarrow \infty$ .

*Proof.* Let  $\chi : \mathbb{R} \rightarrow \mathbb{R}$  be a smooth cutoff function such that  $\chi(t) = 1$  for  $t \leq 1$  and  $\chi(t) = 0$  for  $t \geq 2$ . We define the trace-free symmetric two-tensor  $h$  on  $\mathbb{R}_+^n$  by

$$h_{ab}(x) = \sum_{N=N_0}^{\infty} \chi(4N^2|x-x_N|) 2^{-dN} f(2^N|\bar{x}-x_N|) H_{ab}(x-x_N)$$

where  $x_N = (\frac{1}{N}, 0, \dots, 0) \in \partial\mathbb{R}_+^n$ . Observe that  $h$  is smooth and satisfies  $h_{an}(x) = 0$  for  $x \in \mathbb{R}_+^n$  and  $\partial_n h_{ab}(x) = 0$  for  $x \in \partial\mathbb{R}_+^n$ . If  $N_0$  is sufficiently large, then  $h_{ab}(x) = 0$  for  $|x| \geq \frac{1}{2}$  and  $|h(x)| + |\partial h(x)| + |\partial^2 h(x)| \leq \alpha$  for  $x \in \mathbb{R}_+^n$ , with  $\alpha$  sufficiently small as in Proposition 5.1. Then we define the metric  $g(x) = \exp(h(x))$  for  $x \in \mathbb{R}_+^n$  and the result follows from Proposition 5.1.  $\square$

## Appendix A

In this section we establish some useful identities used in Section 4. They are simple computations which are performed in the Appendix of [6].

**Lemma A-1.** We have  $\int_0^\infty \frac{s^\alpha ds}{(1+s^2)^m} = \frac{2m-\alpha-3}{\alpha+1} \int_0^\infty \frac{s^{\alpha+2} ds}{(1+s^2)^m}$ , for  $\alpha + 3 < 2m$ .

**Proposition A-2.** We have

$$\int_{S^{n-2}} p_k = \frac{r^2}{k(k+n-3)} \int_{S^{n-2}} \Delta p_k$$

for every homogeneous polynomial  $p_k$  of degree  $k$ .

**Corollary A-3.** We have

$$\int_{S^{n-2}} x_i x_j = \frac{\sigma_{n-2}}{n-1},$$

$$\int_{S^{n-2}} x_i x_j x_k x_l = \frac{\sigma_{n-2}}{(n-1)(n+1)} (\delta_{ij} \delta_{kl} + \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk})$$

and

$$\int_{S^{n-2}} x_i x_j x_k x_l x_p x_q = \frac{\sigma_{n-2}}{(n-1)(n+1)(n+3)} (\delta_{ij}\delta_{kl}\delta_{pq} + \delta_{ij}\delta_{kp}\delta_{lq} + \delta_{ij}\delta_{kq}\delta_{lp} \\ + \delta_{ik}\delta_{jl}\delta_{pq} + \delta_{ik}\delta_{jp}\delta_{lq} + \delta_{ik}\delta_{jq}\delta_{lp} \\ + \delta_{il}\delta_{jk}\delta_{pq} + \delta_{il}\delta_{jp}\delta_{kq} + \delta_{il}\delta_{jq}\delta_{kp} \\ + \delta_{ip}\delta_{jk}\delta_{lq} + \delta_{ip}\delta_{jl}\delta_{kq} + \delta_{ip}\delta_{jq}\delta_{kl} \\ + \delta_{iq}\delta_{jk}\delta_{lp} + \delta_{iq}\delta_{jl}\delta_{kp} + \delta_{iq}\delta_{jp}\delta_{kl}).$$

## Appendix B

In this section we establish some results used in Section 4.1. The notations here are the same of that section. In particular, we fix  $a_1 = -1$  and

$$a_0 = \frac{(n+3)(n-6)}{2(n-7)(n-8)} \left\{ 3 + \sqrt{9 - \frac{8(n-7)(n-8)^2(n+7)}{(n+3)(n-6)(n-9)(n-10)}} \right\}.$$

**Proposition B-1.** *We have  $I''(1) < 0$  for  $n \geq 53$ .*

*Proof.* We are going to prove that  $I''(1) < 0$  for  $n \geq 70$ . If  $25 \leq n \leq 69$  the result follows from direct computations. We write

$$a_0 = \frac{(n+3)(n-6)}{2(n-7)(n-8)} \left\{ 3 + \sqrt{9 - \frac{8p_A(n)}{p_B(n)}} \right\},$$

where  $p_A(n) = (n-7)(n-8)^2(n+7)$ ,  $p_B(n) = (n+3)(n-6)(n-9)(n-10)$  and define

$$q_L(n) = p_A(n) - p_B(n) \quad \text{and} \quad q_U(n) = \alpha p_B(n) - p_A(n),$$

where  $\alpha = \frac{31439}{28800}$ .

*Claim.*  $q_L(n) > 0$  for  $n \geq 9$  and  $q_U(n) > 0$  for  $n \geq 70$ .

In order to prove the Claim, first observe that the forth order terms of  $q_L$  cancel out and we have  $q_L(n) = 6n^3 - 114n^2 + 712n - 1516$ . Hence,  $q_L''(n) = 36n - 228 > 0$  for  $n \geq 7$ ,  $q_L(9) = 32$  and  $q_L'(9) = 118$ . Thus,  $q_L(n) > 0$  for  $n \geq 9$ .

Now we observe that

$$q_U(n) = \frac{2639}{28800}n^4 - \frac{115429}{14400}n^3 + \frac{1207877}{9600}n^2 - \frac{282161}{400}n + \frac{218809}{160}.$$

Hence,  $q_U''(n) = \frac{2639}{1200}n - \frac{115439}{2400} > 0$  for  $n \geq 70$ ,  $q_U(70) = \frac{287074}{15}$ ,  $q_U'(70) = \frac{178522037}{7200}$  and  $q_U''(70) = \frac{10910017}{4800}$ . Thus,  $q_U(n) > 0$  for  $n \geq 70$ , proving the Claim.



We assume that  $n \geq 70$ . In particular, we conclude from the Claim that  $\alpha > \frac{p_A(n)}{p_B(n)} > 1$ , which implies

$$\frac{2(n+3)(n-6)}{(n-7)(n-8)} > a_0 > \frac{(n+3)(n-6)}{2(n-7)(n-8)}(3 + \sqrt{9-8\alpha}).$$

Now we use this estimate in

$$I''(1) = \frac{2(n+1)(n-1)}{n-5} \left\{ \frac{a_0^2}{n-6} - \frac{6(n+3)a_0}{(n-8)(n-7)} + \frac{6(n+3)(n+7)}{(n-10)(n-7)(n-9)} \right\}$$

to see that

$$\begin{aligned} \frac{(n-5)I''(1)}{2(n+1)(n-1)} &< \frac{4(n+3)^3(n-6)}{(n-7)^2(n-8)^2} - \frac{3(3 + \sqrt{9-8\alpha})(n+3)^2(n-6)}{(n-7)^2(n-8)^2} \\ &\quad + \frac{6(n+3)(n+7)}{(n-10)(n-7)(n-9)}. \end{aligned}$$

This can be written as

$$I''(1) < \frac{2(n+3)(n+1)(n-1)\gamma(n)}{(n-8)^2(n-10)(n-5)(n-7)^2(n-9)},$$

where

$$\gamma(n) = -(5+3\sqrt{9-8\alpha})(n+3)(n-6)(n-10)(n-9) + 6(n+7)(n-7)(n-8)^2.$$

In order to complete our proof, we will show that  $\gamma(n) < 0$  under our assumption on the dimension. Observe that  $\gamma(n) = -\frac{11}{20}n^4 + \frac{481}{10}n^3 - \frac{15099}{20}n^2 + \frac{21162}{5}n - 8205$ . Hence  $\gamma'''(n) = -\frac{66}{5}n + \frac{1443}{5} < 0$  for  $n \geq 70$ ,  $\gamma(70) = -118392$ ,  $\gamma'(70) = -\frac{744953}{5}$  and  $\gamma''(70) = -\frac{136479}{10}$ . Now the result follows.  $\square$

**Proposition B-2.** *We have  $J(1) < 0$  for  $n \geq 53$ .*

*Proof.* Let us assume that  $n \geq 53$ . We want to show that  $(n+3)\sqrt{9 - \frac{8p_A(n)}{p_B(n)}} - 6 > 0$ , where we are using the polynomials  $p_A$  and  $p_B$  as in the proof of Proposition B-1. We set again  $q_U(n) = \alpha p_B(n) - p_A(n)$  and choose  $\alpha = \frac{7047}{6272}$ .

*Claim.*  $q_U(n) > 0$ .

In order to prove the Claim, first observe that

$$q_U(n) = \frac{775}{6272}n^4 - \frac{27341}{3136}n^3 + \frac{814983}{6272}n^2 - \frac{551233}{784}n + \frac{2063213}{1568}.$$

Hence,  $q_U''(n) = \frac{2325}{784}n - \frac{82023}{1568} > 0$  for  $n \geq 53$ ,  $q_U(53) = \frac{169857}{28}$ ,  $q_U'(53) = \frac{20672955}{1568}$  and  $q_U''(53) = \frac{5182395}{3136}$ . Thus,  $q_U(n) > 0$  for  $n \geq 53$ , proving the Claim.

The Claim implies that  $(n + 3) \sqrt{9 - \frac{8p_A(n)}{p_B(n)}} > (n + 3) \sqrt{9 - 8\alpha}$ , which reduces the problem to prove that

$$(n + 3) \sqrt{9 - 8\alpha} - 6 \geq 0. \quad (\text{B-1})$$

On the other hand, the fact that  $\alpha = \frac{1}{8} \left\{ 9 - \frac{36}{56^2} \right\}$  implies  $\frac{1}{8} \left\{ 9 - \frac{36}{(n+3)^2} \right\} \geq \alpha$ , which is equivalent to the inequality (B-1).  $\square$

## References

- [1] Almaraz, S.: A compactness theorem for scalar-flat metrics on manifolds with boundary. *Calc. Var. Partial Differential Equations*, in press, DOI: 10.1007/s00526-010-0365-8
- [2] Almaraz, S.: An existence theorem of conformal scalar-flat metrics on manifolds with boundary. *Pacific J. Math.* **248**(1), 1-22 (2010)
- [3] Ambrosetti, A., Li, Y., Malchiodi, A.: On the Yamabe problem and the scalar curvature problem under boundary conditions. *Math. Ann.* **322**(4), 667-699 (2002)
- [4] Aubin, T.: Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire. *J. Math. Pures Appl.* **55**, 269-296 (1976)
- [5] Berti, M., Malchiodi, A.: Non-compactness and multiplicity results for the Yamabe problem on  $S^n$ . *J. Funct. Anal.* **180**(1), 210-241 (2001)
- [6] Brendle, S.: Blow-up phenomena for the Yamabe equation. *J. Amer. Math. Soc.* **21**(4), 951-979 (2008)
- [7] Brendle, S., Chen, S.: An existence theorem for the Yamabe problem on manifolds with boundary. *ArXiv:0908.4327v2*
- [8] Brendle, S., Marques, F.: Blow-up phenomena for the Yamabe equation II. *J. Differential Geom.* **81**, 225-250 (2009)
- [9] Chen, S.: Conformal Deformation to Scalar Flat Metrics with Constant Mean Curvature on the Boundary in Higher Dimensions. *ArXiv:0912.1302v2*
- [10] Cherrier, P.: Problèmes de Neumann non linéaires sur les variétés Riemanniennes. *J. Funct. Anal.* **57**, 154-206 (1984)
- [11] Djadli, Z., Malchiodi, A., Ould Ahmedou, M.: Prescribing scalar and boundary mean curvature on the three dimensional half sphere. *J. Geom. Anal.* **13**(2), 255-289 (2003)
- [12] Djadli, Z., Malchiodi, A., Ould Ahmedou, M.: The prescribed boundary mean curvature problem on  $\mathbb{B}^4$ . *J. Differential Equations* **206**(2), 373-398 (2004)

- [13] Druet, O.: From one bubble to several bubbles: the low-dimensional case. *J. Differential Geom.* **63**(3), 399-473 (2003)
- [14] Druet, O.: Compactness for Yamabe metrics in low dimensions. *Int. Math. Res. Not.* **23**, 1143-1191 (2004)
- [15] Escobar, J.: The Yamabe problem on manifolds with boundary. *J. Differential Geom.* **35**, 21-84 (1992)
- [16] Escobar, J.: Conformal deformation of a Riemannian metric to a scalar flat metric with constant mean curvature on the boundary. *Ann. Math.* **136**, 1-50 (1992)
- [17] Escobar, J.: Conformal metrics with prescribed mean curvature on the boundary. *Calc. Var. Partial Differential Equations* **4**, 559-592 (1996)
- [18] Felli, V., Ould Ahmedou, M.: Compactness results in conformal deformations of Riemannian metrics on manifolds with boundaries. *Math. Z.* **244**, 175-210 (2003)
- [19] Felli, V., Ould Ahmedou, M.: A geometric equation with critical nonlinearity on the boundary. *Pacific J. Math.* **218**(1), 75-99 (2005)
- [20] Han, Z., Li, Y.: The Yamabe problem on manifolds with boundary: existence and compactness results. *Duke Math. J.* **99**(3), 489-542 (1999)
- [21] Khuri, M., Marques, F., Schoen, R.: A compactness theorem for the Yamabe problem. *J. Differential Geom.* **81**(1), 143-196 (2009)
- [22] Lee, J., Parker, T.: The Yamabe problem. *Bull. Amer. Math. Soc.* **17**, 37-91 (1987)
- [23] Li, Y., Zhang, M.: Compactness of solutions to the Yamabe problem, II. *Calc. Var. Partial Differential Equations* **24**, 185-237 (2005)
- [24] Li, Y., Zhang, M.: Compactness of solutions to the Yamabe problem, III. *J. Funct. Anal.* **245**(2), 438-474 (2007)
- [25] Li, Y., Zhu, M.: Yamabe type equations on three dimensional Riemannian manifolds. *Commun. Contemp. Math.* **1**(1), 1-50 (1999)
- [26] Marques, F.: A priori estimates for the Yamabe problem in the non-locally conformally flat case. *J. Differential Geom.* **71**, 315-346 (2005)
- [27] Marques, F.: Existence results for the Yamabe problem on manifolds with boundary. *Indiana Univ. Math. J.* **54**(6), 1599-1620 (2005)
- [28] Marques, F.: Conformal deformation to scalar flat metrics with constant mean curvature on the boundary. *Comm. Anal. Geom.* **15**(2), 381-405 (2007)

- [29] Ould Ahmedou, M.: On the prescribed scalar and zero mean curvature on 3-D manifolds with umbilic boundary. *Advanced Nonlinear Studies* **6**, 13-46 (2006)
- [30] Schoen, R.: Conformal deformation of a Riemannian metric to constant scalar curvature. *J. Differential Geom.* **20**, 479-495 (1984)
- [31] Schoen, R.: On the number of constant scalar curvature metrics in a conformal class. *Differential Geometry: A symposium in honor of Manfredo Do Carmo* (H.B.Lawson and K.Tenenblat, eds.), Wiley, 311-320 (1991)
- [32] Schoen, R., Yau, S.-T.: *Lectures on Differential Geometry, Conference Proceedings and Lecture Notes in Geometry and Topology, I.* International Press, Cambridge (1994)
- [33] Schoen, R., Zhang, D.: Prescribed scalar curvature on the n-sphere. *Calc. Var. Partial Differential Equations* **4**(1), 1-25 (1996)
- [34] Trudinger, N.: Remarks concerning the conformal deformation of a Riemannian structure on compact manifolds. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)* **22**, 265-274 (1968)
- [35] Yamabe, H.: On a deformation of Riemannian structures on compact manifolds. *Osaka Math. J.* **12**, 21-37 (1960)

INSTITUTO DE MATEMÁTICA  
 UNIVERSIDADE FEDERAL FLUMINENSE  
 NITERÓI - RJ, BRAZIL  
 E-mail addresses: [almaraz@vm.uff.br](mailto:almaraz@vm.uff.br), [almaraz@impa.br](mailto:almaraz@impa.br)